

ACCEPTED  
 Apr. 16, 1970  
 UNDER FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENT ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 1388-25

*With  
 Comments*

**DEADLY**  
 POISONED GRAIN  
 for  
 GROUND SQUIRRELS  
 and MICE



**Directions—Ground Squirrels**

When putting this poison out use teaspoonful to a tablespoonful of bait distributed near the entrance of burrows and along trails made by the squirrels. This bait has been found effective on the Oregon and Douglas ground squirrels. The best time for poisoning these squirrels is early spring around March or when they become active, as they are hungry and take the bait more readily.

BIRDS FEEDING ON TREATED AREAS MAY BE KILLED

**Directions for House Mice**

Place rounded teaspoon of bait at 8 to 10 foot intervals where mice feed, water or travel. Place in locations not accessible to children, pets and domestic animals. Collect and destroy all uneaten baits when through.

BAIT SHOULD BE PLACED IN PROTECTIVE BAIT BOXES

USDA Reg. 1388-25

Packaged by  
 U.S. RODENT DESTROYER CO.

Division of  
**KORINEK REMEDY CO.**  
 STAYTON, OREGON 97269

**DEADLY**  
 POISONED GRAIN


**FOR USE AGAINST  
 GROUND SQUIRRELS  
 and MICE**

**DEADLY**  
 POISONED GRAIN FOR  
 GROUND SQUIRRELS & MICE

**DEADLY**  
 POISONED GRAIN  
 for  
 GROUND SQUIRRELS  
 and MICE



**"DANGER"** Keep Out of Reach of Children

**POISON**  **WARNING:** Convulsive Poison! Do not contaminate feed and foodstuffs. Keep away from children and domestic animals.

**ANTIDOTE:** If less than ten minutes has passed since the poison was taken, give a tablespoonful of salt in a glass of warm water. Have victim lie down in a quiet, darkened room and keep warm. Call a Physician immediately.

ACTIVE INGREDIENTS  
 Strychnine as Alkaloid 9.5%  
 INERT INGREDIENTS 90.5%  
 NET WEIGHT 1 LB

**DEADLY**  
 POISONED GRAIN

**DEADLY**  
 POISONED GRAIN FOR  
 GROUND SQUIRRELS & MICE

**FOR USE  
 AGAINST  
 GROUND  
 SQUIRRELS  
 and MICE**

**FOR USE AGAINST  
 GROUND SQUIRRELS  
 and MICE**

ACCEPTED  
 Apr 16 1970  
 FOR ECONOMIC PROTECTIVE FIELD  
 UNDER NO 1388-25

**DEADLY**  
 POISONED GRAIN  
 for  
 GROUND SQUIRRELS  
 and MICE



**Directions—Ground Squirrels**  
 When putting this poison out use teaspoonful to a tablespoonful of bait distributed near the entrance of burrows and along trails made by the squirrels. This bait has been found effective on the Oregon and Douglas ground squirrels. The best time for poisoning these squirrels is early spring around March or when they become active, as they are hungry and take the bait more readily.

BIRDS FEEDING ON TREATED AREAS MAY BE KILLED

**Directions for House Mice**  
 Place rounded teaspoon of bait at 8 to 10 foot intervals where mice feed, water or travel. Place in locations not accessible to children, pets and domestic animals. Collect and destroy all uneaten baits when through.

BAIT SHOULD BE PLACED IN PROTECTIVE BAIT BOXES

USDA Reg. 1388-25  
 Packaged by  
 U.S. RODENT DESTROYER CO.  
 Division of  
 KORINEK REMEDY CO.  
 STAYTON, OREGON 97163

**DEADLY**  
 POISONED GRAIN

FOR USE AGAINST  
 GROUND SQUIRRELS  
 and MICE

**DEADLY**  
 POISONED GRAIN FOR  
 GROUND SQUIRRELS & MICE

**DEADLY**  
 POISONED GRAIN  
 for  
 GROUND SQUIRRELS  
 and MICE



"DANGER" Keep Out of Reach of Children

**POISON**  **WARNING:**  
 This poison is highly toxic to humans and animals. It is fatal if swallowed. It is also highly toxic to birds and fish. Do not use in areas where these animals are present.

**ANTIDOTE:** If swallowed, induce vomiting immediately. If inhaled, move to fresh air. If on skin, wash with soap and water. If on clothing, remove and wash separately.

**DEADLY**  
 POISONED GRAIN FOR  
 GROUND SQUIRRELS & MICE

FOR USE AGAINST  
 GROUND  
 SQUIRRELS  
 and MICE

**DEADLY**  
 POISONED GRAIN

FOR USE AGAINST  
 GROUND SQUIRRELS  
 and MICE