

2, 4-D WEED KILLER

Kills Dandelion, Goldenrod, Ragweed, Broad and Narrow Leaf Plantain, Thistle, Dock, Wild Morning-glory, Shepherd's Purse, Wild Carrot, Mallow and many other broad leaved weeds without injury to most grasses.

Active Ingredients:

Alkanolamine Salts (of the ethanol and isopropanol series)

of 2, 4-Dichlorophenoxyacetic acid\* 10%

Inert Ingredients: 90%

\*Equivalent to 6.2% 2,4-Dichlorophenoxyacetic acid

DIRECTIONS

2, 4-D WEED KILLER is highly soluble in water and adaptable to preparation sprays. Always use sufficient water to insure distribution of 2, 4-D WEED KILLER over area being treated, i.e., about 125 gallons per acre. Spray while weeds are growing actively and after leaves are fully developed, but prior to bud development. Since the response of weeds in different areas to 2, 4-D compounds is so greatly influenced by environmental conditions, users should consult State Agricultural Experimental Station or Extension Service Weed Specialists.

On pastures, lawns, golf courses, parks, playgrounds and other turf areas: for control of broad leaf weeds such as plantain, dandelion and chickweed, and shallow rooted weeds such as yellow mustard and ragweed, use 16 pints per acre. This is equivalent to about 1 pound of 2, 4-D acid equivalent per acre, or 6 ounces of 2, 4-D WEED KILLER in 3 gallons of water, for every 1,000 square feet.

For weeds which are difficult to control such as yarrow, goldenrod and wild onions, use about 20 pints per acre. This is equivalent to about 7 1/2 ounces of 2, 4-D WEED KILLER in 3 gallons of water, for every 1,000 square feet.


In vacant lots and air fields, on roadsides, along fence rows and railroad right-of-ways, around farm buildings and other waste areas; for control of deep rooted perennial weeds such as Canada thistle, bindweed, poison ivy, and sow thistle, use about 24 pints per acre. This is equivalent to about 9 ounces of 2, 4-D WEED KILLER in 3 gallons of water for every 1,000 square feet.

**IMPORTANT:** Do not apply 2, 4-D WEED KILLER directly to, or otherwise permit it to contact, vegetables, flowers, fruit trees, ornamentals, cotton or other desirable plants which are sensitive to 2, 4-D and similar materials. Do not permit spray mist containing it to drift onto them, since even minute quantities of the spray might cause severe injury. Coarse sprays are less likely to drift.

Do not use on lawns of creeping grasses, except for spot spraying, nor on seeded lawns until the grass has become well established. Most legumes are usually damaged or killed. Excessive amounts of 2, 4-D in the soil may temporarily inhibit seed germination or plant growth. Do not use weed control equipment for applying insecticides or fungicides. Do not store near fertilizers, seeds, insecticides or fungicides. Do not graze dairy animals within 7 days after treatment. Do not contaminate water used for irrigation or domestic purposes.

Do not re-use empty container. Destroy by burying in a safe place.

CAUTION (18 pt.): Keep out of reach of children (12 pt.)  
May cause skin irritation. Avoid contact with eyes, skin or clothing.  
wash off promptly with plenty of water.



U.S.D.A. Registration No. 1416-14

Net Vol. Gals.

Manufactured by CHURCHILL CHEMICAL COMPANY, Galesburg, Illinois