TRIPLE PLAY

Detergent - Germiciae		TR: My
High Potency Disinfectant-Cleaner for		th
Hospital and Other Environmental Sanitation	n Programs	(F Se
A_0 , A_0 . Use Dilution: 100 to 1		CA
Tuberculocidal at 80 to 1		ca
Cleans, Disinfects and Deodorizes in one Operation. Non-Selective. Kills many Gram Positive and Gram Negative Organisms. Kills most common pathogenic bacteria and fungi, including anti-biotic resistant Staph. <u>Biodegradable</u> .		FI me co
For Cleaning, Disinfecting and Deodorizing all type of surfative Walls, and Fixtures in Hospitals, Sick Rooms, Contagious Dis Schools, Office Buildings, Factories and Institutions.	sease Areas, Wash Rooms,	U.
Tested and approved for use on conductive	e hospital floors.	
Especially designed for use where soil and bacter	rial contamination are high.	77
Active ingredients: Vegetable Oil Soap Potassium-o-benzyl-p-chlorophenate Sodium alkyl (C9) benzene sulfonate Ethylene diamine tetracetic acid, ter Inert ingredients: 83.32% Water Tetrasodium pyrosphosphate Sodium Nitrite Perfume Dye	trasodium salt ACCEPTED AUG-0.1967	

. A March Ol 1. 1. Ch. 1 C

Directions

FOR CLEANING, DISINFECTING AND DEODORIZING IN ONE OPERATION: Make solution of 1 part TRIPLE PLAY to 100 parts warm water, this is approximately 1 1/4 fluid ounces per gallon. Apply this solution to contaminated surface and scrub lightly with mop or brush. Let it remain on surface for several minutes, then remove and rinse with fresh water. For extra dirty areas use a little stronger solution.

IN TUBERCULOSIS SANATORIUMS or wherever virulent concentrations of Mycobacterium tuberculosis are suspected, use solution containing 1 part of TRIPLE PLAY to 80 parts of water that is, about 1 2/3 ounces per gallon.

TRIPLE PLAY passes A.O.A.C. Use Dilution. Confirmation Test against Staphylococcus aureus and Salmonella choleraesuis at 100 to 1 dilution. Accordingly, it is accepted for hospital and other environmental sanitation programs.

IPLE PLAY also passes the most recently adopted A.O.A.C. test procedure against cobacterium Tuberculosis at 80 to 1, after incubation for 90 days -- in ree different media.

ront Panel) CAUTION (18 pt.): Kccp out of reach of children (12 pt.) e side panel for additional precautions.

UTION: Harmful if taken internally. Keep out of reach of children. May use skin irritation. Avoid contact with skin. Avoid splashing in eyes.

RST AID: In case of skin contact, flush with plenty of water; for eyes, get dical attention. If swallowed, give raw egg white, milk, mucilage or ornstarch paste and follow with emetic of mustard. Call a physician.

S.D.A. Registration No. 1416-17

Gals. Net Vol.

Manufactured by CHURCHILL CHEMICAL COMPANY, Galesburg, Ill.

