

KEM KO 6 WEED-KILLER

A total non-selective weed killer for application on non-cropland areas where bare ground is desired. Ready to use.

ACTIVE INGREDIENTS:

Refined Aliphatic Hydrocarbons	56.10%
Isocetyl Ester of 2,4-Dichlorophenoxyacetic Acid*	1.09%
Bromacil (5-Bromo-3-sec-butyl-6-methyluracil)	0.61%
Pentachlorophenol	0.80%
Other Chlorophenols	0.09%
INERT INGREDIENTS:	1.31%

Total 100.00%

*Equivalent to 4.22% of 2,4-Dichlorophenoxyacetic Acid.

WARNING

KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED OR ABSORBED THROUGH THE SKIN. VAPOR HARMFUL. CAUSES SKIN IRRITATION. If swallowed, do not induce vomiting. Call physician immediately. Do not breathe vapor or mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. In case of contact with skin, wash with soap and water. In case of contact with eyes, immediately flush with water for at least 15 minutes and get medical attention. Remove and wash contaminated clothing before re-use. Do not contaminate feed or foodstuffs.

This product is toxic to fish and wild life. Keep out of lakes, streams, ponds. Do not contaminate any body of water. Wildlife in treated areas may be harmed.

For use only on non-crop lands. Do not apply directly or permit this product to come in contact with any desired plant or vegetable. Do not permit spray mists to drift to other areas. Use a coarse spray. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply to areas underlaid by roots of desired plants.

Thoroughly clean spray equipment before using for any other purposes. Reserve spray equipment for this purpose only. Do not store near fertilizers, insecticides, seeds, or any plant treating materials. Use this product only for the purpose intended. Do not re-use this container. Rinse empty container with water and puncture to prevent re-use. Do not burn container. Bury with all rinsings and spillage in non-agriculture area away from all water supplies.

FLAMMABLE LIQUID: Do not store, pour, or spray near open flame or heat.

DIRECTIONS:

Apply any time of year except when ground is frozen. Best results are obtained by treating after predominant species are well established in late spring or early summer. The degree and duration of results will depend upon the amount applied, type of soil, amount of rainfall and other conditions. User should consult agricultural Experiment Station or Extension Service weed specialists as to use of this product in their locality or consult the sales representative.

General Weed Control: Apply 2 to 5 gallons per 1,000 square feet (80 to 200 gals. per acre) on non-cropland areas as parking lots, fence rows, loading ramps, railroad sidings, storage yards, around buildings and industrial plant sites to control most weeds for an extended period of time.

Annual Weeds and Grasses, as foxtail, ryegrass, wild oats, crabgrass, cheatgrass, broomsedge, ragweed, lambquarters and turkey mullein, apply 1 to 2 gallons per 1,000 square feet (40 to 80 gallons per acre).

Perennial Weeds and Grasses, as smoothbrome, Canada bluegrass, purple top, quackgrass, broomsedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot, apply 2 to 3 gallons per 1,000 square feet (80 to 150 gallons per acre).

For Hard to Kill Perennial Weeds, as Johnsongrass, salt grass and horsetail, apply 3 to 5 gallons per 1,000 square feet (120 to 220 gallons per acre).

USDA Registration No. 1481-181

Net Contents

Gallons

SOLD BY

DEITELBACH CHEMICAL CORPORATION

4181 Peachtree Road, N. W.

Atlanta, Georgia 30319

ACCEPTED
7-31-69
1921-161
FUNGICIDE
N. J. DEITELBACH CHEMICAL CORP. PHILADELPHIA, PA.

A total non-selective weed killer for application on
ground is desired. Ready to use.

ACTIVE INGREDIENTS:

Refined Aliphatic Hydrocarbons
Isooctyl Ester of 2,4-Dichlorophenoxyacetic acid
Bromacil (5-Bromo-3-sec-butyl-6-methyluracil)
Pentachlorophenol
Other Chlorophenols

INERT INGREDIENTS:

Equivalent to 4.52% of 2,4-Dichlorophenoxyacetic acid

WARNING

KEEP OUT OF THE REACH OF CHILDREN

HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN. HARMFUL. CAUSES SKIN IRRITATION. If swallowed, call physician immediately. Do not breathe vapor, skin, or on clothing. Wash thoroughly after handling with soap and water. In case of contact with eyes, flush with water for at least 15 minutes and get medical attention. Wash clothing before re-use. Do not contaminate food or feed. This product is toxic to fish and wild life. Keep out of water. Do not contaminate any body of water. Wildlife in treated areas may be affected. For use only on non-crop lands. Do not apply directly to any desired plant or vegetable. Do not apply to other areas. Use a coarse spray. Vapors from this product may injure plants in the immediate vicinity. Do not apply to desirable plants.

Thoroughly clean spray equipment before using for other spray equipment for this purpose only. Do not use on seeds, or any plant treating materials. Use this product only on non-crop lands. Do not re-use this container. Rinse empty container before re-use. Do not burn container. Bury with agricultural waste in an agriculture area away from all water supplies.

FLAMMABLE LIQUID: Do not store, pour, or spray near open flame.

DIRECTIONS:

Apply any time of year except when ground is frozen. Treat after moderate frosts are well established. The degree and duration of results will depend on soil amount of rainfall and other conditions. Experiment Station or Extension Service weed specialist in their locality or consult the sales representative.

General Weed Control: Apply 2 to 3 gallons per 1,000 square feet (10 to 15 gallons per acre) on non-crop lands such as parking lots, tennis courts, sidewalks, storage yards, around buildings and industrial areas for an extended period of time.

Annual Weeds and Grasses, as foxtail, cypergrass, bromeseed, crabgrass, and turkey in 1,000 square feet (40 to 50 gallons per acre).

Perennial Weeds and Grasses, as smoothbrome, Canada grass, barnyardgrass, aster, dandelion, dog fennel, goldfennel, apply 2 to 3 gallons per 1,000 square feet (80 to 100 gallons per acre).

For Hard to Kill Weeds such as Johnsongrass, apply 3 to 5 gallons per 1,000 square feet (130 to 220 gallons per acre).

USEDA Registration No. 1421-161

SC 10 37