

ACCEPTED  
 Jan 26, 1969 # (1421-183)  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 6754-18

Comments

**MALATHION**  
**FLY BAIT**

WITH D. D. V. P.  
**Attracts and Kills**  
**Resistant House Flies**

ACTIVE INGREDIENTS:

Malathion (O, O-Dimethyl Dithiophosphate of Diethylmercaptosuccinate	1.0%
O, O Dimethyl 2, 2 Dichloro Vinyl Phosphate	.5%
INERT INGREDIENTS	98.5%

**CAUTION: READ CAUTION STATEMENT**  
**KEEP OUT OF REACH OF CHILDREN**

**DIRECTIONS**

Apply Fly Bait daily while flies are abundant. Sprinkle thinly in strips on floor and other places where flies gather, but not where it will contaminate feed or food. Usually, about 1 oz. (2 tbsp.) is needed for each 500 sq. ft. depending on the number of flies. If they are extremely numerous, several times this quantity may be necessary. Fly Bait is especially recommended for use in barns, stables, pens, feed rooms, poultry houses, kennels, manure piles, garbage cans, and other outside areas. In dairy barns, treat the feed room floor and the barn floor, but do not feed the floor sweepings to livestock. Repeat as needed. In poultry houses where poultry is in cages, use Fly Bait on walkways in and around the building, including feed rooms and egg houses. Where poultry is on ground or litter, apply only outside building or in yard: where poultry cannot feed on Fly Bait.

To kill fly maggots, apply Fly Bait evenly and thoroughly on manure at the rate of 1 to 1 1/2 lbs. per 100 sq. ft. (2 to 2 1/2 oz. per sq. yd.). If in small mounds, apply on sides as well as top. Repeat in 5 to 7 days, and afterwards as needed.

TO MAKE A LIQUID BAIT: Dissolve in water at rate of 1 pound per gallon. Spray solution in coarse spray and apply to point of run-off. Apply daily until fly population is under control. Repeat applications as necessary. Do not use as a spray in dairy barns.

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid skin contact. Wash thoroughly with soap and water after using. Keep container out of reach of children and animals. Avoid contamination of food, feed, and utensils used for food or feed. Do not use inside homes or where milk and other foods are stored or processed. Repeated or careless use may cause Cholinesterase inhibition. Atropine is antidotal.

U.S.D.A. REG. NO. 6754-18  
 NET WEIGHT POUND

ACCEPTED WITH COMMENTS

# FUNDAL<sup>®</sup> 4EC

## MITICIDE/INSECTICIDE/OVICIDE

### GENERAL INFORMATION

FUNDAL 4EC is a water emulsifiable miticide insecticide ovicide for the control of certain insects and mites of fruit trees, cotton and cole crops.

### ENVIRONMENTAL HAZARD

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### PRECAUTIONS

It is not necessary to use a spreader sticker with FUNDAL 4EC. However, if one is used it is recommended that the tolerance of the crop to such mixture be determined by first testing it on a small scale. Do not spray shortly before rainfall. Prepare fresh sprays daily. Do not allow prepared solutions to stand overnight. Flush tank thoroughly with water daily after use. Do not use FUNDAL 4EC in alkaline solutions such as those containing lime sulfur or Bordeaux. Do not allow animals to graze in areas treated with FUNDAL 4EC. Do not feed crop residues

E.P.A. Reg. No. 2139-100

### ACTIVE INGREDIENTS:

Chlordimeform*	48.5%
Aromatic Petroleum Distillates	43.4%
INERT INGREDIENTS:	8.1%
Total	100.0%

Contains 4 lbs. Chlordimeform per gallon  
N (4-Chloro o-tolyl) N,N-dimethylformamide

### WARNING

### KEEP OUT OF REACH OF CHILDREN

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles when handling. Avoid inhaling vapors or spray mist. Wash with soap and water after handling and before eating or smoking. Wear clean clothes and wash before reusing. Do not contaminate food or feedstuffs.

In case of contact, thoroughly clean skin and clothing. For eyes, flush immediately with water and get medical attention.

**Do not reuse this container.  
Destroy when empty.**

U.S. PAT. NO. 3,178,407 (5-10-72)

## NET CONTENTS

### STORAGE AND HANDLING INSTRUCTIONS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not store at temperatures below 32° F. Crystals may form if stored under very cold conditions. If this should occur place container in a warm room (50-80° F) and agitate frequently for several days by rolling and shaking the container until crystals disappear.

### IMPORTANT

### READ BEFORE PURCHASE OR USE

Purchase or use of this product shall constitute acceptance of the following terms of sale. NOR AM Agricultural Products, Inc. (NOR AM) warrants that this product conforms to the chemical descriptions on this label and is reasonably fit for the purposes specified hereon. By way of limitation but not in derogation of the directions and statements printed on this label, Seller makes no warranty and no representation and NOR AM makes no further warranty or representation, expressed or implied, including but not limited to any warranty as to MERCHANTABILITY, and no agent of NOR AM is authorized to do so except in writing, with a specific reference to the foregoing warranty. Since neither Seller nor NOR AM has any control over the handling, storage, use or conditions of use of this product, Buyer or User assumes all risk and liability arising from handling, storage or use of this product under abnormal conditions or not in strict compliance with the directions and precautionary statements hereon. All claims related to the purchase, handling, storage or use of this product shall be barred unless written notice is given to NOR AM within 30 days after the date of purchase.