METHYL DODECYLBENZYL TRIMETHYL AMMONIUM CHLORIDE 4% METHYL DODECYLXYLENE BIS (TRIMETHYL AMMONIUM CHLORIDE 1%

INERT INGREDIENTS INCLUDES SPECIAL DETERGENTS

HARMFUL IF SWALLOWED
MAY CAUSE SKIN BURNS OR DAMAGE TO THE EYES
DO NOT GET IN EYES OR ON SKIN. IN CASE OF
CONTACT WITH EYES FLUSH WITH PLENTY OF
WATER FOR 15 MINUTES AND GET MEDICAL ATTENTION.
AVOID INHALATION OF DUST.

DETERGENT SANITIZER FOR WASHING EGGS

CONTAINS
HYAMINE
A.QUARTERNARY
AMMONIUM
GERMICIDE

DANGER

KEEP OUT OF REACH OF CHASEE SIDE PANEL FOR PRECA

U.S.D.A. REG. NO. 1430-13

ACTIVIT INGREDIENTS ACTIVIT INGREDIENTS ACTIVIT INGREDIENTS ACTIVIT INGREDIENTS ACTIVIT INGREDIENTS SOCIUM Metasificate, Pentahydrate Sodium Carbonate 1079 Sodium Carbonate INGREDIENTS A0% INGREDIENTS TOTAL CLEEDES SPECIAL DETERGENTS 100%

The use of CO-OP BRAND DETERGENT SANITIZER for washing soiled eggs will help reduce spoilage of fresh and storage eggs due to bacterial thereon. This product is an excellent sanitizing detergent for washing feed hoppers, water founts, and other equipment in poultry houses and egg handling rooms.

OF BOTH HOSBALLOWED

Made in U.S.A. NET WEIGHT LBS.

CO-OP BRAND

DETERGENT SANITIZER FOR WASHING EGGS

CONTAINS

A QUARTERNARY

AMMONIUM

GERMICIDE

20% 40% 60% 100% bacterial makes. This rooms.

T WEIGHT LBS.

CO-OP BRAND

DETERGENT
SANITIZER
FOR WASHING EGGS

CONTAINS
A QUARTERNARY
AMMONIUM
GERMICIDE

1. Keep pens clean — wash only soiled eggs.

DIRECTIONS FOR WASHING EGGS:

2. Dissolve 1 oz. (1 measure) of Detergent Sanitizer in each 3 gallons of water (110°-130° F.). Maintain wash solution at even temperature. Very dirty eggs may require stronger solution.

3. For spray-type machines follow manufacturer's directions.

4. For hand washing or immersion-type machine, immerse eggs for 3 minutes. Remove from solution, allow to drain. Do not wash more than 5 dozen eggs per gallon of solution. Change solution oftener if it becomes dirty.

5. If separate rinse is used, add 1 oz. Chlorinated Detergent per 5 gallons of water. Maintain rinse at temperature slightly greater than wash solution.

6. Dry eggs rapidly and store in cool, damp room.

DIRECTIONS FOR GENERAL CLEANING AND SANITIZING:

- 1. Prepare hot water solution containing 1 oz. of detergent sanitizer per 2 gallions of water.
- 2. Wash feed hoppers, water founts and other equipment thoroughly. In case of disease outbreak, apply fresh solution after cleaning.
- 3. Wash equipment, walls, and floors of egg handling rooms to REDUCE BACTERIAL CONTAMINATION AND MALESTANDER.

Distributed by INTER-COUNTY FARMERS CO-OP ASSN., INC. Woodridge, N. Y.