


WILDLIFE

-This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and, upon repeated feedings, causes fatal hemorrhage in rats.

Place baits in locations not accessible to children, pets, and domestic animals or in tamper-proof bait boxes.

INTER
r-County Farmers Cooperative Association Inc.
WOODRIDGE, NEW YORK



**READY
TO USE**

CO-OP BRAND Bait

KILLS RATS AND MICE

ACTIVE INGREDIENTS:

N'-(2-quinoxaliny) sulfanilamide (sulfquinoxaline) 0.025%

INERT INGREDIENTS99.950%


***Trade mark of the Wisconsin Alumni Research Foundation**

U.S.D.A. No. 1430-53

CAUTION—KEEP OUT OF THE REACH OF CHILDREN

187 TYPE ¹ See left panel for additional cautions.

NET WEIGHT 3 LBS.



BRAND Bait

RATS AND MICE

..... 0.025%
..... 0.025%
..... 99.950%

Research Foundation

S.D.A. No. 1430-53

OUT OF THE REACH OF CHILDREN

For additional cautions.

NET WEIGHT 3 LBS.

READY
TO USE

1. GENERAL DIRECTIONS

- Place baits in locations where rats and mice feed, water, or travel.
- Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.
- Contaminated or spoiled baits should be replaced promptly.
- Baiting should continue until all signs of feeding have stopped.

2. SPECIAL DIRECTIONS

- Where labeling has directions for using fresh meat or fish as bait, also state, "Dogs and cats will actively search out baits of this type. Therefore, baits must be placed so as to prevent feeding by these animals".
- The following statement may be included, but is not mandatory; "Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed."

3. DIRECTIONS FOR RAT CONTROL (Sketches to illustrate are desirable)

- Four ounces of bait per station should be the minimum amount used.
- Bait should be placed in locations not accessible to children, pets, ^{WILDLIFE} and domestic animals, or in tamper-proof bait boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.

4. DIRECTIONS FOR MOUSE CONTROL (Sketches to illustrate are desirable)

- Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8 = 12 foot intervals.
- Bait should be placed in locations not accessible to children; pets, ^{WILDLIFE} and domestic animals, or in tamper-proof bait boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.