

1439-60  
ACCEPTED  
MAR 1 1968

[Faint, mostly illegible text from a document, possibly a letter or report, with some lines of text visible.]

BEST DOCUMENT AVAILABLE

POTASSIUM CYANATE  
100% PURE  
NET WT. 5 LBS.

ACTIVE INGREDIENTS . . . . .  
Potassium Cyanate 95%  
INERT INGREDIENTS 5%

INSTRUCTIONS:

To control young crabgrass seedlings early in the  
year, two applications of Potassium Cyanate s  
tablespoon per gallon water, over 100 square fe  
should be one week apart. Test sprinkler can f  
get an idea of coverage. For mature crabgrass,  
application at double the strength (2 tablespoo  
enough, but repeat if necessary. Do not apply  
when desirable grasses are started or dormant,  
winter. Five gallons treat 1200 square feet.

Potassium Cyanate spray should be applied when  
crabgrass seedlings, crabgrass or chickweed will di  
cause no injury to desirable turf. Any  
by the first cut following treatment.  
to be used fresh. When used as directed, t  
residue in soil, breaks down into harm

Keep out of hands of children.

We do not warrant or guarantee that the application  
of this product of any kind, express or implied, or  
we do not assume all risks for use of this product  
in any way.