

BEST DOCUMENT AVAILABLE

BEST DOCUMENT

DIRECTIONS FOR USE: This product may be applied by either low pressure, low volume equipment or by conventional high pressure sprayer. It may be added to either water or solvents in mixing application. The time of application is determined largely by the stage of growth of the weeds, but when applied as a selective spray the stage of growth of the crop plant must be considered.

SMALL GRAIN CROPS: (Oats, Barley, Wheat, Rye): Apply 1 pint per acre on susceptible weeds as wild pigweed and ragweed and 1-1-1/2 pints per acre on perennial weeds such as bindweed. Late application any time after grain plants are 1" to 2" tall except blossoming stage shall be avoided.

CORN: For annual weeds employ 1/2 - 3/4 pint per acre; perennial weeds will require 1 - 1-1/2 pints per acre. Avoid application of solution to growing point of corn if possible.

PLANT: Susceptible weed plants require 1/3 - 1/4 pint per acre. Apply when plants are 4-6" high. Be especially careful in following dosage for this crop.

PASTURES: Apply 1-2 pints per acre on annual weeds and 1-3 quarts per acre for perennials.

FRUIT TREES, ROADSIDES, DITCH BANKS: Apply 1-2 quarts per acre as soon as weeds appear and 2 quarts per acre if woody plants such as poison oak, sumac or other brush are present. Avoid contamination of weeds used for irrigation.

UNPLANTED AREAS: Apply 1 pint per acre on annuals and 1 quart per acre on perennials.

WIND AND SOLE PAINMENT: Apply 1-2 pints per acre depending on infestation and season of year. Avoid use on bent grass and clover.

Use above quantities to liquid carrier at time of application. Quantity of spray solution will depend on equipment but in general 3-5 gallons in low volume sprayers and 175-150 gallons in high volume equipment is sufficient per acre. Low pressures of 15-50 lbs. are most desirable. Wind, at areas growing weeds are most susceptible. Two to four weeks may pass before kill is evident; perennial weeds may require retreatment of areas infested.

CAUTION: Avoid spray drift to susceptible plants as 2,4-D will injure cotton, beans, peas, ornamentals, etc. (Cotton sprays are less likely to drift). In addition, vapors from this product may injure susceptible plants in the immediate vicinity. DO NOT use the same spray equipment for applying other materials to plants as injury will result. DO NOT store near fertilizers, seeds, insecticides, or fungicides. Avoid contact with skin, eyes or clothing.

DO NOT USE ON NEWLY SEED-BED WEEDS.

GENERAL WEEDER

2,4-D WEED KILLER

LOW VOLUME

Contains 4 lbs. 2,4-D acid Equivalent per gallon

ACTIVE INGREDIENTS:
Isobutyl ester of 2,4-Dichlorophenoxyacetic Acid... 70.5%

INERT INGREDIENTS:..... 29.5%

TOTAL:..... 100.0%

*Equivalent to 2,4-Dichlorophenoxyacetic acid..... 46.84%

NOTICE: The agents concern only. They are conditions of day control, nor does it a warranty, guarantee concerning the final description Corporation's manner for all type of loss handling, etc. accordance with liability requirements and

NON-WARRANTY

DO NOT USE ON NEWLY SEED-BED WEEDS

BEST DOCUMENT AVAILABLE

COMMENTS

HEM WEED

2,4-D WEED KILLER

LOW VOLATILE

(1)

NOTICE: The statements made on this label, or by any of our agents concerning this material, are given for information only. They are believed to be true and accurate, but because conditions of use which are of critical importance are beyond our control, Chemical Insecticide Corporation does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither Chemical Insecticide Corporation nor the seller shall be held responsible in any manner for any personal injury, or property damage or other type of loss resulting to the buyer or to any person from the handling, storage, or use of this material, whether or not in accordance with directions. The buyer assumes all risk and liability resulting from such handling, storage or use and accepts and uses this material on these conditions.

NOT-GUARANTEE
Seller gives no warranty of any kind, expressed or implied, concerning the use of this product. Buyer assumes all risks of use or handling whether in accordance with directions or not.

Contains 4 lbs. 2,4-D acid Equivalent per gallon

ACTIVE INGREDIENTS:	
Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid	79.5%
INERT INGREDIENTS:..... 20.5%	
TOTAL.....100.0%	

Equivalent to 2,4-Dichlorophenoxyacetic acid..... 46.8%

Do not use on sensitive plants & fruit trees
SAME NAME AS FERTICOR.

ACCEPTED
1439-101
FEB 7 - 1968
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND ROUPICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. SUBJECT TO ATTACHED COMMENTS.

Amendments:

- (1) WARNING: KEEP OUT OF THE REACH OF CHILDREN
"See side panel for additional required caution statements."
- (2) CHEMICAL INSECTICIDE CORPORATION
30 Whitman Avenue
Metuchen, New Jersey
Lalle Glade, Florida.....Presque Isle, Maine
- (3) "U.S.D.A." Reg. No. 1439-101
- (4) PASTURES: Do not graze treated areas by dairy animals within 7 days after treatment.
- (5) SMALL GRAIN CROPS:
 - a. "Do not forage or graze grain fields for 2 weeks after treatment with 2,4-D."
 - b. "Do not use treated straw for livestock feed."

ACCEPTED
Feb. 7, 1968
UNDER THE FEDERAL INSECTICIDE ACT
FOR REGISTERED PRODUCTS
UNDER NO. 1439-101

Comment