CONTROL NO.

DIRECTIONS FOR USE

First Read Warning and Caution on this label

GENERAL. [illegible]

[illegible] AND [illegible] PLANT CONTROL May be applied [illegible] ground sprayers or by airplane in a mixture of water and [illegible] diesel oil Make up spray by adding CHEM VEX to the [illegible] keep agitating during spraying Apply when [illegible] to growth at parts Annual follow up applications may be [illegible]

Ground Use Ground Sprayer [illegible] quart per 100 gallon water per acre [illegible] quarts [illegible] adequate for [illegible]

Airplane Application [illegible] quarts in [illegible] gallons of water per acre [illegible] gallon of diesel oil alone

Stump Application Apply when new sprouts have made good growth and before frost tops growth

Spot Treatment For backpack applications [illegible] pint in 3 gallons of water spray to wet foliage

Species Controlled Very effective [illegible] following

Oak [illegible] oak, black [illegible] shingle oak and Northern oak

Maple [illegible] red sugar and big leaf maple

Mesquite [illegible] and Grease Wood Wild Blackberry Bramble Yucca Prickly Pear [illegible] effective on spruce, ash and honeylocust than ordinary brush killer [illegible] 2,4-D and 2,4,5-T

Herbaceous Broadleaves [illegible] Weeds [illegible] Croton Weed, Broom Weed, Lamsquarters, [illegible] Mexican Weed Pokeberry, Ragweed Sunflower Plantain

AQUATIC WEEDS. Submerged Aquatics Effective for the control of certain submerged aquatic weeds [illegible] Water Milfoil Myriophyllum spp Fanwort Cabomba [illegible] Waterweed Helodea spp Water Najad Najas spp Flatsedge [illegible] Anacharis spp Water Starwort [illegible] 1 to 2 parts per million ppm [illegible] to 5 quarts per [illegible] or early summer when weeds are actively [illegible] using sufficient water [illegible] of spray [illegible]

[illegible] control of [illegible] Naiad [illegible] deep water [illegible] and Pickerel weed [illegible] of CHEM VEX [illegible] gallons [illegible]

CHEM - SECT BRAND

CHEM VEX

A Low Volatile Brush and Weed Killer

(Contains 4 pounds of silvex per gallon)

For use on roadsides, rights-of-way, fence rows, ditch banks and forest management areas. For selective broad-leaved weed control in rice and on established turf, and for selective control of certain aquatic weeds in lakes and ponds.

CAUTION Keep out of reach of children.

See side panel for additional required caution statements

ACTIVE INGREDIENT:

Isooctyl ester of silvex

(2-(2,4,5-trichlorophenoxy) propionic acid)* 65.1%

INERT INGREDIENTS 34.9%

*Equivalent to 46.5% by weight of silvex or not less than 4 lbs of silvex per gallon at 68 F

CAUTION [illegible] Avoid breathing spray mist May cause skin irritation [illegible] contact with eyes, skin and clothing In case of contact [illegible] with plenty of water wash skin with soap and water Do not [illegible] with feed seeds or foodstuffs Do not contaminate water [illegible] for domestic or recreational purposes Thoroughly clean [illegible] equipment before [illegible] for other purposes

USDA REG. NO 1439 174

[illegible] Proper identification of treated [illegible] essential to avoid dis[illegible] intended for the control of Duckwe[illegible] spp [illegible] algae [illegible] use [illegible] cultural [illegible] comply with local and state reg[illegible] duction of chemicals into waters Do not apply copper sulfate for [illegible] Treat only ponds or lakes with [illegible] may reduce the effectiveness of [illegible] practical to reduce water flow [illegible] aquatic animals, some fish kill m[illegible] coves and where uniform applicati[illegible]

WEED CONTROL IN TURF: For sele[illegible] and other established turf areas [illegible] leaved plantain buckhorn, this mouse-eared chickweed, dandeli[illegible] clover and black medic. Apply 2 [illegible] per acre (approximately 1 fluid [illegible] 1,000 square feet Best applied [illegible] if used in very hot weather may [illegible] apply to newly sown turf nor[illegible] Bermuda or St Augustine

WEED CONTROL IN RICE where [illegible] weed, Mexican weed curly, and broadleaf [illegible] use [illegible] to 3 [illegible] water for even distribution For adequate [illegible] Apply 4 to 8 weeks [illegible] treat before 7 to [illegible] weeks af[illegible] above water and leaves are sta[illegible] usually safer than before [illegible] Extension Service for specific inca[illegible]

WARNING: Be sure [illegible] use [illegible] regulations Do Not apply CHE[illegible] drift to [illegible] crops [illegible] peanuts [illegible] sweet potatoes [illegible] ornamentals [illegible]

[illegible]

CHEMICAL INSECTICIDE CORPORATION

30 WHITMAN AVENUE, METUCHEN, NEW JERSEY U. S. A.

PRESQUE ISLE, MAINE

BELLE GLADE, FLORIDA