

3/27/68

Formal No. _____

CHEM. SECT. LABEL

PARATHION 15% EMULSION POWDER

1439-129 Comments

1. PRECAUTIONS:

(1)

1.1. First Aid: If swallowed, inhaled, or absorbed through skin, wash with water. Do not get in eyes or on skin. Wear natural rubber gloves, goggles, and a mask or respirator of a type passed by the U.S. Dept. of Labor for parathion protection. Keep all unprotected persons out of vicinity where there may be danger of drift. Vapors may be inhaled until drifting, insecticide, and volatile residues may be on hands, contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with water before eating or smoking. Wash all contaminated clothing in hot water before re-use. Burn or destroy, never reuse containers. Do not use spillage with alkali. Stay out of smoke or fumes.

1.2. First Aid: If swallowed, give a tablespoonful of salt in a glass of water. Induce vomit fluid is clear. Have victim lie down and keep quiet. **ATROPINE IMMEDIATELY!** ATROPINE IS ANTIDOTAL. Keep an emergency supply of atropine. **CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.** SEE PHYSICIAN'S NOTE BELOW.

1.3. First Aid: Remove patient from contaminated atmosphere, keep him under observation. Don't wait for doctor, give atropine (2 tablets) at once if symptoms of blurred vision or pin-point pupils.

1.4. First Aid: Remove contaminated clothing and scrub skin with soap and water. Don't wait for doctor, give atropine (2 tablets) at once if symptoms of blurred vision or pin-point pupils.

1.5. First Aid: If swallowed, make patient vomit by giving warm salty or soapy water. Don't wait for doctor, give atropine (2 tablets) at once if symptoms of blurred vision or pin-point pupils. Have victim lie down and keep quiet.

1.6. First Aid: NOTHING BY MOUTH TO AN UNCONSCIOUS PATIENT.

1.7. First Aid: Warning symptoms include weakness, headache, chills, blurred vision, pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea.

1.8. First Aid: Give atropine preferably by injection, grains 1 to 3 mg. at once and parenterally or orally every hour as required until pupils dilate, secretions dry, and clear chest by coughing. Use artificial respiration if breathing becomes convulsed or stops.

1.9. First Aid: Observe patient continuously as he may have repeated exposure to vapors. Patients may, without warning, have prolonged susceptibility to the effects of any cholinesterase inhibitor. No further exposure until cholinesterase regeneration has been determined by blood test.

2. Use: 10% EMULSION: VEGETA: 1 lb. per 100 gal. of water. Beans, corn, cotton, eggplants, and tomatoes, use 1 lb. per 100 gal. of water. Apply at 10 day intervals. Do not apply later than 30 days after blossoming whichever is earlier. For aphids, use 1 lb. per 100 gal. of water. For scale, use 15 lbs. per 100 gal. of water. For various and celery, use either 1 lb. per 100 gal. of water or 100 lbs. per acre. DO NOT APPLY WITHIN 30 DAYS OF HARVEST. For control of pear aphid, use 1% dust when aphids appear or after blossoming.

3. Warnings: KEEP OUT OF THE REACH OF CHILDREN. See label panel for additional precautionary statements.

ACCEPTED WITH COMMENTS