

DIRECTIONS FOR USE

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

COCKROACHES — Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

ANTS — Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

SILVERFISH, FIREBRATS AND PAPERLICE — Apply to localized areas infested by these insects.

BROWN DOG TICKS — Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS AND SPIDERS — Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention around window frames.

FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES AND SAND-FLIES (outdoors only) — Do not treat lawns, herbaceous plants, vines, shrubs, and trees. Contamination of ornamental ponds may be harmful to fish. Birds feeding on treated area may be harmed. Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

CAUTION

May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Should poisoning occur, get prompt medical aid.
To physician — Atropine sulfate is antidotal.

DO NOT REUSE EMPTY CONTAINER. DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

"DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME."

TOMIC



ACTIVE INGREDIENTS: 100%

0 - Isopropoxyphenyl Methylcarbamate	1%
2 - Butoxyethanol	9%
Petroleum Distillate	90%

® BAYGON is the registered trademark of
Farbenfabriken BAYER A.G., Chemagro
Corporation Licensee.

KILLS

Cockroaches, Ants, Brown Dog Ticks, Centipedes, Clover Mites, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Millipedes, Mosquitoes, Paperlice, Punkies, Sandflies, Scorpions, Silverfish, Sowbugs and Spiders.



CAUTION: KEEP OUT OF THE REACH OF CHILDREN

(See Side Panel for Additional Cautions)

NET CONTENTS 1 GALLON (128 FL. OZ.)

DIRECTION FOR USE (continued)

Insect Control in Research Facilities:

The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

RESTRICTIONS

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in dining areas while food is exposed.

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

This product is toxic to birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

For Industrial and Institutional Use

Residual Spray is a highly effective ready to use formulation for control of the insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This product drives insects from their hiding places, gives rapid kill and residual activity up to 30 days.

Ca. Reg. No. 01440-50001 AA
E.P.A. Est. No. 1440-CA-1
EPA Reg. No. 1440-CA-7

Manufactured By

TOMIC INSECTICIDE COMPANY

241 So. Mednik Avenue
Los Angeles, California 90022

1440-7

2/17/77

PS 181