

NOTE TO PHYSICIAN

Antidote — administer atropine sulfate in large doses. TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

POST TREATED AREA

Consult your State Agricultural Service or Experiment Station regarding posting treated areas.

STOP - READ LABEL - BEFORE USING

Agro-Chemical Brand

CHEMICAL DISTRIBUTORS

O B A



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

EPA REG. NO. 1526-208 AA.

EPA EST. NO. 1526-AZ-1

PARATHION 2

ACTIVE INGREDIENTS:

Parathion: O,O-diethyl O-p-nitrophenyl phosphorothioate

2%

INERT INGREDIENTS:

98%

100%

POISON

See Side Panels for Antidote & Precautions



DANGER

Keep Out of Reach of Children

NOT FOR HOME USE.

Out of Reach of Children. Precautions on this label




NOTICE

Small text notice regarding product use and safety.

DIRECTION

CROP	PEST	APPL.
Alfalfa	Armyworms, Alfalfa Caterpillars, Aphids.	40 lbs. per acre.
Broccoli, Brussel Sprouts, Cauliflower	Aphids, Cabbage Looper.	25 lbs. per acre.
Cabbage	Aphids, Cabbage Looper.	25 lbs. per acre.
Carrots	Aphids	50 lbs. per acre.
Cotton	Cotton Leaf Perforator, Cotton Fleahopper, Stink Bugs.	60 lbs. per acre.
Garlic	Thrips	30 lbs. per acre.
Lettuce (head)	Armyworms, Aphids, Cabbage Loopers.	25 lbs. per acre.
Lettuce (Bibb leaf)	Armyworms, Cabbage Looper.	12 lbs. per acre.
	Aphids	25 lbs. per acre.
Melons	Aphids, Leafhoppers	25 lbs. per acre.
Onions	Thrips	40 lbs. per acre.
Potatoes	Aphids, Fleabeetles, Grasshoppers, Potato Psyllid.	50 lbs. per acre.
Radishes	Aphids	25 lbs. per acre.
Safflower	Aphids	25 lbs. per acre.
Sugar Beets	Aphids, Armyworms	40 lbs. per acre.
Sweet Potatoes	Aphids, Leafhoppers.	25 lbs. per acre.
Tomatoes	Aphids, Armyworms, Cabbage Loopers.	25 lbs. per acre.



TO READ LABEL BEFORE USING!

Agro-Chemical Brand

CHEMICAL DISTRIBUTORS

DBA



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

EPA REG. NO. 1526-208 AA.
EPA EST. NO. 1526-AZ-1

PARATHION 2

ACTIVE INGREDIENTS:


Parathion: O,O-diethyl O-p-nitrophenyl
phosphorothioate

2%

INERT INGREDIENTS:

98%

100%



NOTICE

THIS PRODUCT IS A PESTICIDE. IT IS TO BE USED ONLY ON THE CROPS AND PLANTS SPECIFICALLY LISTED ON THE LABEL. IT IS TO BE USED ONLY IN THE MANNER AND AT THE RATES SPECIFICALLY LISTED ON THE LABEL. IT IS TO BE USED ONLY BY PERSONS WHO HAVE BEEN TRAINED AND CERTIFIED IN THE PROPER USE OF PESTICIDES.

DIRECTIONS FOR USE

CROP	PEST	APPLICATION	LIMITATIONS
Alfalfa	Armyworms, Alfalfa Caterpillars, Aphids.	40 lbs. per acre. In California 20 lbs. per acre.	Do not apply within 15 days of harvest.
Broccoli, Brussel Sprouts, Cauliflower	Aphids, Cabbage Looper.	25 lbs. per acre.	Do not apply within 7 days of harvest.
Cabbage	Aphids, Cabbage Looper.	25 lbs. per acre.	Do not apply within 10 days of harvest.
Carrots	Aphids.	50 lbs. per acre.	Do not apply within 15 days of harvest. Do not use treated tops for food or feed.
Cotton	Cotton Leaf Perforator, Cotton Fleahopper, Stink Bugs.	60 lbs. per acre.	Workers entering fields within 24 hours after application should wear protective clothing. Do not hand pick or harvest within 7 days of application.
Garlic	Thrips	30 lbs. per acre.	Do not apply within 15 days of harvest.
Lettuce (head)	Armyworms, Aphids, Cabbage Loopers.	25 lbs. per acre.	Do not apply within 7 days of harvest.
Lettuce (Bibb leaf)	Armyworms, Cabbage Looper.	12 lbs. per acre.	Do not apply within 14 days of harvest.
	Aphids	25 lbs. per acre.	Do not apply within 21 days of harvest.
Melons	Aphids, Leafhoppers	25 lbs. per acre.	Do not apply within 7 days of harvest.
Onions	Thrips	40 lbs. per acre.	Do not apply within 15 days of harvest.
Potatoes	Aphids, Fleabeetles, Grasshoppers, Potato Psyllid.	50 lbs. per acre.	Do not apply within 5 days of harvest.
Radishes	Aphids	25 lbs. per acre.	Do not apply within 15 days of harvest.
Safflower	Aphids	25 lbs. per acre.	Do not apply after flowering.
Sugar Beets	Aphids, Armyworms	40 lbs. per acre.	Do not apply within 15 days of harvest.
Sweet Potatoes	Aphids, Leafhoppers	25 lbs. per acre.	Do not apply within 15 days of harvest.
Tomatoes	Aphids, Armyworms, Cabbage Loopers	25 lbs. per acre.	Do not apply within 10 days of harvest.

