A READY-TO-USE, NON-SELECTIVE VEGETATION KILLER FOR BARE-GROUND WEED CONTROL AROUND FENCES, GUARD RAILS, HIGHWAY MEDIANS, PIPELINES, FIRE HYDRANTS, PUMPS, RAILROAD TRACKS, STORAGE AREAS, FUEL TANKS AND MACHINERY


IMPORTANT: READ ENTIRE DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY DISCLAIMER BEFORE USING THIS PRODUCT.



E"A Hen. Mo. 14-9-55

Bullen

CAUTION: Keep Out of Reach of Children Harmful if swallowed. Avoid contact with eyes and skin. If swallowed do not induce vomiting. Call a physician immediately. In case of skin contact wash with soap and water. Avoid inhalation of spray mist... Avoid contamination of food and feed. Do not contaminate water used for irrigation or demestic purposes...Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label . . . Do not reuse container. Destroy when empty...DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Store at temperatures above 32° F.



VEGETATION KILLER

WARRANTY DISCLAIMER:

Since weather and other conditions under which this product may be used will vary. the Seller makes no warranty of any kind, express or implied, concerning the use of this product. The user assumes all risks of use or handling not in accordance with directions or suggestions

PRECAUTIONS

Care must be taken to keep this herbicide away from desirable flower beds. lawns and shrubs. Do not apply over roots of valuable trees or woody ornamental plants. Only apply herbicide outside the dripline of trees. Avoid spray drift to desirable plants.

Do not reuse empty container. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place av ay from water supplies.

BULLEN CHEMICAL CO., FOLCROFT, PA. 19032

ACTIVE INGREDIENTS

William Regions . By

Prometon: 2.4-bis (isopropylamino)-	
6-methoxy-s-triazine	1.5%
Petroleum distillate	 94.0%
INERT INGREDIENTS	 4.5%
Total .	 100.00%

NET: 5 U.S. GALLONS

1459 PA 1

DIRECTIONS FOR USE:

Use 2½ to 3 gallons undiluted weed-killer per 1000 sq. ft. of soil surface (100-200 gal. per acre). Spray uniformly and thoroughly wet all foliage to insure contact action. Best results are obtained if used when weeds are just beginning to sprout and are growing rapidly.

Where vegetation is tall and dense, use 5 gals, per 1000 sq. ft. (200 gal. per acre). For control of hard-to-kill perennial weeds and grasses such as Johnson grass, bindweed and wild carrott, use 8 to 12 gals, per 1000 sq. ft.

Although it has considerable activity through foliage contact, much of this weed killers activity is through the roots and, therefore, adequate rainfall or watering is necessary.

Under conditions where the rainfall is high and the growing season long, a second application may be necessary to maintain a weed-free soil surface.

CASTION: Keep Out Of Reach Of Shildren. SEE SIDE PANELS FOR ADDITIONAL CAUTION STATEMENTS