

FOR THE RECORD OF THE COURT
OF THE DISTRICT OF COLUMBIA
IN THE MATTER OF THE ESTATE OF

[illegible]

VALCO

CHEMICALS

Manufactured by **VALCO CHEMICALS DIVISION** Harling
VALLEY CO-OP OIL MILL

RESTRICTED USE PESTICIDE

Use only by Certified Applicators or persons under their direct supervision, and only as directed by the Certified Applicator's certification.

VALCO

CHEMICALS DIVISION

MAY 12 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1467-61

METHYL PARATHION 7.5 EC

(O,O-Dimethyl O-p-Nitrophenyl) Phosphorothioate	70.55%
Isopropyl Alcohol	3.71%
Aromatic Petroleum Derivative Solvents	18.57%
Other Ingredients	7.17%
TOTAL	100.00%

1467-TX-01
1467-61

CONTAINS 7½ POUNDS METHYL PARATHION PER GALLON

STATEMENT OF PRACTICAL TREATMENT

ANTIDOTE-CONSULT PHYSICIAN FOR EMERGENCY SUPPLY
 IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING
 If victim is awake (conscious), make him vomit quickly. Have victim drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting by mouth to an unconscious person. Have victim lie down and keep quiet. Get patient immediately from the area where methyl parathion is present. Get medical attention immediately. Flush skin with plenty of water while removing contaminated clothing and shoes. Flush eyes for at least 15 minutes with plenty of water. Help patient by breathing fresh air. Get medical attention.

KEEP OUT OF THE REACH OF CHILDREN

DANGER-POISON

Study precautions, poisoning symptoms, First Aid Treatment and Antidote given on left panel of this label

VALCO CHEMICALS DIVISION Harlingen, Texas 78550
 VALLEY CO-OP OIL MILL

DIRECTIONS FOR USE

Read and follow Federal Law to use this pesticide in a manner inconsistent with directions on the label.
TO PREVENT PERSONAL INJURY AND POSSIBLE FATALITIES
 Workers entering treated fields within 48 hours must wear protective clothing. Treated areas should not be reentered until drifting insecticide and volatile residues have dissipated. To avoid excessive residues of methyl parathion on food or forage crops, always observe the statements found on the label before harvest. Methyl parathion may be applied because this material is poisonous by skin contact, inhalation or swallowing. It should not be used in such a manner or under weather conditions that will permit drift of the spray onto areas not intended to be sprayed. Do not apply with hand equipment. Add the concentrate to the spray tank while filling with water and mix thoroughly by means of continual tank agitation during filling and spraying operation. For application by ground equipment add the desired amount of concentrate to sufficient water to apply at least 3 gallons of spray mix per acre. For application by aircraft add the desired amount of concentrate to sufficient water to apply 1 to 3 gallons of spray mix per acre. For best results thoroughly cover all surfaces to be treated with spray.
COTTON For control of thrips use 1.8 to 1.4 pint per acre. For cotton leafworms use 1.8 to 3.8 pint per acre. For grasshoppers and fall armyworms use 1.4 pint per acre. For spider mites, do not control all species; use 1.4 to 3.8 pint per acre. For control of leafhoppers, Lygus and other mites use 1.8 pint to 1 pint per acre. For boll weevils, aphids and garden webworms use 1.4 to 1.2 pint per acre. For false chinch bugs use 1.2 pint per acre. For cutworms use 1.2 to 1 pint per acre. For bollworms, salt marsh caterpillars, armyworms, cotton leaf perforators and stink bugs use 1 pint per acre. Application should be made at 4 to 5 days before all cotton bolls are opened. At above dosages application may be made up to the day before harvest. Do not use with mechanical pickers. Workers entering treated fields within 48 hours must wear protective clothing and NOT APPLY WITHIN 5 DAYS OF HANDPICKING COTTON.

TANK MIXTURE RATES FOR COTTON POUNDS ACTIVE INGREDIENT/ACRE

Methyl parathion	EPN	Toxaphene	Chlor-dimethform	Permethrin
1/2-3/4	1/2-3/4	—	1/8	—
1/2-1	—	1-2	—	—
1	—	2	1/8	—
1/2-1	—	—	1/8	—
1/2-1	—	—	—	0.05-0.1

BEANS Dry and Green For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs and mites use 1.8 to 3.8 pint per acre. For leaf beetles, surface feeding and climbing cutworms, and stink bugs use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 21 DAYS OF HARVEST.
BEETS Red For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs use 1.8 to 3.8 pint per acre. DO NOT APPLY 15 DAYS OF HARVEST. If logs are to be used for food or feed, DO NOT APPLY WITHIN 21 DAYS OF HARVEST.
CABBAGE COLE CROPS Broccoli, Sprouts, Cauliflower, Kohlrabi, Rutabagas, COLLARDS, KALE AND MUSTARD GREENS For control of aphids, armyworms, flea beetles, leafhoppers and mites use 1.8 to 3.8 pint per acre. For stink bugs use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 21 DAYS OF HARVEST.
CARROTS For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites use 1.8 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST. Do not use treated carrot tops for food or feed.
CELERY For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
CUCUMBERS For control of aphids and two-spotted mites use 1.4 pint per acre. Do not apply before or after DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
CORN For control of armyworms, climbing cutworms, corn leaf aphids, corn rootworm adults, fall armyworms, and stink bugs use 1.4 pint per acre. For grasshoppers use 1.2 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
LETTUCE For control of aphids, armyworms, flea beetles, leafhoppers and mites use 1.8 pint per acre. DO NOT USE WITHIN 21 DAYS OF HARVEST.
ONIONS For control of thrips use 1.4 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
PEPPERS For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites use 1.8 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
POTATOES For control of aphids, armyworms, false chinch bugs, flea beetles, leafhoppers, mites and stink bugs use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.
SPLASH For control of aphids, armyworms, flea beetles, leafhoppers, mites, seed corn maggots, and other insects use 1.2 to 1 pint per acre. DO NOT APPLY WITHIN 21 DAYS OF HARVEST.
SWEET POTATOES For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites use 3.4 pint per acre. DO NOT APPLY WITHIN 5 DAYS OF HARVEST.
TOMATOES For control of aphids, armyworms, flea beetles, leafhoppers, mites and psyllids use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST.
TURNIPS For control of aphids, armyworms, flea beetles, leafhoppers, leaf miners and mites use 1.8 to 3.8 pint per acre. DO NOT APPLY WITHIN 15 DAYS OF HARVEST. If logs are to be used for food or feed, DO NOT APPLY CLOSER THAN 21 DAYS OF HARVEST.

CONDITIONS OF SALE

Read directions carefully. Timing and method of application, weather and crop conditions, along with other chemicals not specifically recommended, and other influencing factors in the use of this product are the responsibility of the user. Buyer assumes all risks of use, storage or handling of this material in strict accordance with the directions given on the label.

Label No. 8008-154