

TREFLAN® E.C. (E.P.A. REG. NO. 1471-35)
TREFLAN® M.T.F.™ (E.P.A. REG. NO. 1471-116)
TREFLAN® 5™ (E.P.A. REG. NO. 1471-120)

TREFLAN/Command® and TREFLAN/Command/Lexone® or
Sencor® combinations for Weed Control in Soybeans
(Do Not Use In California)

TREFLAN may be tank-mixed with Command or Command plus Lexone or
Sencor for weed control in soybeans.

TREFLAN/Command Tank-Mix

TREFLAN/Command will control the following weeds:

Grasses

Annual bluegrass
Barnyardgrass
 (Watergrass)
Brachiaria
 (Signalgrass)
Bromegrass
 (Cheatgrass)
 (Downy brome)
Cheat
 (Chess)
Crabgrass
 (Large crabgrass)
 (Smooth crabgrass)
Foxtail
 (Bottlegrass)
 (Bristlegrass)
 (Giant foxtail)
 (Green foxtail)
 (Foxtail millet)
 (Pigeongrass)
 (Robust foxtail)
 (Yellow foxtail)

Goosegrass
 (Silver crabgrass)
 (Silvergrass)
 (Wiregrass)
 (Yardgrass)
Johnsongrass
 (from seed)
Junglerice
Texas panicum
 (Buffalograss)
 (Coloradograss)
Sandbur
 (Burgrass)
Sprangletop
Stinkgrass
 (Lovegrass)
Woolly cupgrass

(Florida purslane)
(Mexican clover)
(Pusley)
Goosefoot
Knotweed
Kochia
(Fireweed)
(Mexican fireweed)
Lambsquarters

(Redroot)
(Rough pigweed)
(Spiny pigweed)
Purslane
Russian thistle
(Tumbleweed)
Stinging nettle
(Nettle)
Velvetleaf
(Buttonweed)

TREFLAN/Command will provide partial control or suppression of the following weeds:

Jimsonweed
Morningglory, annual
Prickly sida
(Teaweed)

Ragweed, common
Smartweed, Pennsylvania
Venice mallow

Control of these weeds may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

Application Directions

Tank-mixed TREFLAN/Command must be applied as a preplant incorporated treatment. Apply TREFLAN/Command up to 3 weeks prior to planting.

Do not apply aerially or through irrigation equipment.

Apply TREFLAN/Command using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10-15 gallons per acre. Use nozzles suitable for broadcast boom application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays.

Incorporation Directions

TREFLAN/Command must be incorporated immediately after application. TREFLAN/Command should be thoroughly and uniformly mixed into the top 2 to 3 inches of the final seed bed. Application to overly

152****/EC**48/LABELS/US/P0789/180

moist or wet soils will increase the potential for off-site movement of Command vapors and may result in poor soil incorporation and unsatisfactory weed control. These directions must be followed to reduce the potential for off-site movement of Command vapors and potential injury to desirable vegetation including adjacent crops, trees and ornamentals.

Follow other soil preparation application and incorporation procedures described on the TREFLAN label.

Broadcast Rates Per Acre

<u>Soil Texture</u>	<u>TREFLAN</u>		<u>Command 4 E.C. (Pints)</u>
	<u>E.C./M.T.F. (Pints)</u>	<u>or TREFLAN 5 (Pints)</u>	
Coarse	1	0.8	3/4
Medium	1 1/2	1.2	1 1/8
Fine	2	1.6	1 1/2

TREFLAN/Command/Lexone or Sencor Tank-Mix

TREFLAN/Command/Lexone or Sencor controls those weeds listed for TREFLAN/Command plus these additional weeds:

Jimsonweed	Sesbania, hemp
Prickly sida	Smartweed, Pennsylvania
(Teaweed)	Spotted spurge
Ragweed, common	Venice mallow ¹
	Wild mustard ¹

TREFLAN/Command/Lexone or Sencor also provides partial control or suppression of common cocklebur, annual morningglory and giant ragweed. Control of these weeds may be erratic ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

Applications Directions

Tank-mixed TREFLAN/Command/Lexone or Sencor must be applied as a preplant incorporated treatment. Apply TREFLAN/Command/Lexone or Sencor up to 2 weeks prior to planting.

Do not apply aerially or through irrigation equipment.

Apply TREFLAN/Command/Lexone or Sencor using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10-15 gallons per acre. Use nozzles suitable for broadcast boom application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays.

Incorporation Directions

TREFLAN/Command/Lexone or Sencor must be incorporated immediately after application. TREFLAN/Command/Lexone or Sencor should be thoroughly and uniformly mixed into the top 2 to 3 inches of the final seed bed. Application to overly moist or wet soils will also increase the potential for off-site movement of Command vapors and may result in poor soil incorporation and unsatisfactory weed control. These directions must be followed to reduce the potential for off-site movement of Command vapors and potential injury to desirable vegetation including adjacent crops, trees and ornamentals.

Follow other preparation application and incorporation procedures described on the TREFLAN label.

Broadcast Rates Per Acre

Soil Texture	TREFLAN		Command 4 E.C. (Pints)	Lexone 4L or Sencor 4F (Pints)	Lexone Dry Flowable or Sencor Dry Flowable (Pounds)
	E.C./M.T.F. (Pints)	or TREFLAN 5 (Pints)			
Coarse	1	0.8	1/2	1/3-1/2 ¹	1/4-1/3 ¹
Medium	1 1/2	1.2	3/4	1/2-3/4	1/3-1/2
Fine	2	1.6	1 1/8	3/4-1	1/2-2/3

1. Use the higher rate in the rate range in areas where weed populations are dense, for control of Venice mallow and wild mustard and for best control of common cocklebur, annual morningglory and giant ragweed.

Precautions

Special Precaution

Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of adjacent crops, trees and ornamental plants which is usually temporary in nature but can result in permanent injury or death of the plants if the exposure is excessive. Prior to making applications of this product, read and strictly follow all precautions and application instructions on the Command label.

Precaution

Read the TREFLAN, Lexone or Sencor labels carefully for additional directions, precautions and limitations before applying these products.

TREFLAN® (trifluralin, Elanco)

Command® (FMC)

Lexone® (metribuzin, DuPont)

Sencor® (metribuzin, Bayer)

Elanco Products Company
A Division of Eli Lilly and Company
Indianapolis, Indiana 46285