

TREFLAN® EC (EPA REG. NO. 1471-35)  
TREFLAN® M.T.F.™ (EPA REG. NO. 1471-116)  
TREFLAN® 5 (EPA REG. NO. 1471-120)  
TREFLAN® TR-10® (EPA REG. NO. 1471-143)

Treflan for Foxtail Control in Spring Seeded Barley Used as a Cover Crop in the Acreage Conservation Reserve Program

Treflan may be applied in the spring as a preplant soil incorporated treatment for foxtail control in spring seeded barley grown on land used in acreage conservation reserve programs.

Apply and incorporate Treflan following recommended procedures in the product labels. Treflan E.C. and Treflan M.T.F. should be applied at the rate of one (1) pint per acre on coarse-textured soils and one and one-half (1½) pints per acre on medium- and fine-textured soils. Treflan 5 should be applied at the rate of eight-tenths (0.8) pints per acre on coarse-textured soils and one and two-tenths (1.2) pints per acre on medium- and fine-textured soils. Treflan TR-10 should be applied at the rate of five (5) pounds per acre on coarse-textured soils and seven and one-half (7½) pounds per acre on medium- and fine-textured soils. Do not exceed this rate or crop injury may occur.

#### Planting Directions

Barley should be seeded approximately two (2) inches deep.

#### Precaution

Use of this practice may result in a slight stand reduction. Do not graze or feed forage from treated fields to livestock. For all land uses, follow current USDA Acreage Conservation Reserve Program guidelines.

Read the Treflan labels for additional directions, precautions and limitations before use.

Treflan® E.C. (trifluralin, Elanco)  
Treflan® M.T.F.™ (trifluralin, Elanco)  
Treflan® 5 (trifluralin, Elanco)  
Treflan® TR-10® (trifluralin, Elanco)

Elanco Products Company  
A Division of Eli Lilly and Company  
Indianapolis, Indiana 46285

TREFLAN® EC (EPA REG. NO. 1471-35)  
TREFLAN® M.T.F.™ (EPA REG. NO. 1471-116)  
TREFLAN® 5 (EPA REG. NO. 1471-120)  
TREFLAN® TR-10® (EPA REG. NO. 1471-143)

Treflan for Foxtail Control in Spring Seeded Barley Used as a Cover Crop in the Acreage Conservation Reserve Program

Treflan may be applied in the spring as a preplant soil incorporated treatment for foxtail control in spring seeded barley grown on land used in acreage conservation reserve programs.

Apply and incorporate Treflan following recommended procedures in the product labels. Treflan E.C. and Treflan M.T.F. should be applied at the rate of one (1) pint per acre on coarse-textured soils and one and one-half (1½) pints per acre on medium- and fine-textured soils. Treflan 5 should be applied at the rate of eight-tenths (0.8) pints per acre on coarse-textured soils and one and two-tenths (1.2) pints per acre on medium- and fine-textured soils. Treflan TR-10 should be applied at the rate of five (5) pounds per acre on coarse-textured soils and seven and one-half (7½) pounds per acre on medium- and fine-textured soils. Do not exceed this rate or crop injury may occur.

#### Planting Directions

Barley should be seeded approximately two (2) inches deep.

#### Precaution

Use of this practice may result in a slight stand reduction. Do not graze or feed forage from treated fields to livestock. For all land uses, follow current USDA Acreage Conservation Reserve Program guidelines.

Read the Treflan labels for additional directions, precautions and limitations before use.

Treflan® E.C. (trifluralin, Elanco)  
Treflan® M.T.F.™ (trifluralin, Elanco)  
Treflan® 5 (trifluralin, Elanco)  
Treflan® TR-10® (trifluralin, Elanco)

Elanco Products Company  
A Division of Eli Lilly and Company  
Indianapolis, Indiana 46285

TREFLAN® EC (EPA REG. NO. 1471-35)  
TREFLAN® M.T.F.™ (EPA REG. NO. 1471-116)  
TREFLAN® 5 (EPA REG. NO. 1471-120)  
TREFLAN® TR-10® (EPA REG. NO. 1471-143)

Treflan for Foxtail Control in Spring Seeded Barley Used as a Cover Crop in the Acreage Conservation Reserve Program

Treflan may be applied in the spring as a preplant soil incorporated treatment for foxtail control in spring seeded barley grown on land used in acreage conservation reserve programs.

Apply and incorporate Treflan following recommended procedures in the product labels. Treflan E.C. and Treflan M.T.F. should be applied at the rate of one (1) pint per acre on coarse-textured soils and one and one-half (1½) pints per acre on medium- and fine-textured soils. Treflan 5 should be applied at the rate of eight-tenths (0.8) pints per acre on coarse-textured soils and one and two-tenths (1.2) pints per acre on medium- and fine-textured soils. Treflan TR-10 should be applied at the rate of five (5) pounds per acre on coarse-textured soils and seven and one-half (7½) pounds per acre on medium- and fine-textured soils. Do not exceed this rate or crop injury may occur.

#### Planting Directions

Barley should be seeded approximately two (2) inches deep.

#### Precaution

Use of this practice may result in a slight stand reduction. Do not graze or feed forage from treated fields to livestock. For all land uses, follow current USDA Acreage Conservation Reserve Program guidelines.

Read the Treflan labels for additional directions, precautions and limitations before use.

Treflan® E.C. (trifluralin, Elanco)  
Treflan® M.T.F.™ (trifluralin, Elanco)  
Treflan® 5 (trifluralin, Elanco)  
Treflan® TR-10® (trifluralin, Elanco)

Elanco Products Company  
A Division of Eli Lilly and Company  
Indianapolis, Indiana 46285

152\*\*\*\*/EC\*\*48/LABELS/US/F0789/247

TREFLAN® EC (EPA REG. NO. 1471-35)  
TREFLAN® M.T.F.™ (EPA REG. NO. 1471-116)  
TREFLAN® 5 (EPA REG. NO. 1471-120)  
TREFLAN® TR-10® (EPA REG. NO. 1471-143)

Treflan for Foxtail Control in Spring Seeded Barley Used as a Cover Crop in the Acreage Conservation Reserve Program

Treflan may be applied in the spring as a preplant soil incorporated treatment for foxtail control in spring seeded barley grown on land used in acreage conservation reserve programs.

Apply and incorporate Treflan following recommended procedures in the product labels. Treflan E.C. and Treflan M.T.F. should be applied at the rate of one (1) pint per acre on coarse-textured soils and one and one-half (1½) pints per acre on medium- and fine-textured soils. Treflan 5 should be applied at the rate of eight-tenths (0.8) pints per acre on coarse-textured soils and one and two-tenths (1.2) pints per acre on medium- and fine-textured soils. Treflan TR-10 should be applied at the rate of five (5) pounds per acre on coarse-textured soils and seven and one-half (7½) pounds per acre on medium- and fine-textured soils. Do not exceed this rate or crop injury may occur.

#### Planting Directions

Barley should be seeded approximately two (2) inches deep.

#### Precaution

Use of this practice may result in a slight stand reduction. Do not graze or feed forage from treated fields to livestock. For all land uses, follow current USDA Acreage Conservation Reserve Program guidelines.

Read the Treflan labels for additional directions, precautions and limitations before use.

Treflan® E.C. (trifluralin, Elanco)  
Treflan® M.T.F.™ (trifluralin, Elanco)  
Treflan® 5 (trifluralin, Elanco)  
Treflan® TR-10® (trifluralin, Elanco)

Elanco Products Company  
A Division of Eli Lilly and Company  
Indianapolis, Indiana 46285

152\*\*\*\*/EC\*\*48/LABELS/US/F0789/247