

SURFLAN® 75W (EPA Reg. No. 1471-96)
SURFLAN® A.S. (EPA Reg. No. 1471-112)

SURFLAN FOR FALL OR EARLY SPRING APPLICATION
FOR NO-TILL SOYBEANS

SURFLAN A.S. and SURFLAN 75W may be applied in the fall or early spring directly over undisturbed stubble from the previous crop. SURFLAN application can begin anytime following harvest after October 15. Spring applications may be made up to four (4) weeks prior to planting but not later than April 15.

The soil should not be tilled between the SURFLAN application and soybean planting.

Early applied SURFLAN effectively controls these annual grasses and broadleaf weeds:

Grasses

Barnyardgrass

(Watergrass)

Brachiaria

(Signalgrass)

Browntop panicum¹

Crabgrasses

(Large crabgrass)

(Smooth crabgrass)

Crowfootgrass

Fall panicum

(Spreading panicgrass)

Foxtails

(Bottlegrass)

(Bristlegrass)

(Giant foxtail)

(Green foxtail)

(Pigeongrass)

(Robust foxtail)

(Yellow foxtail)

Goosegrass

(Silver crabgrass)

(Wiregrass)

(Yardgrass)

Johnsongrass

(Seedling only)

Texas panicum¹

(Buffalograss)

(Coloradograss)

ACCEPTED

NOV 28 1984

Under Federal Insecticide,
Fungicide, and Rodenticide Act,
exempted from the pesticide
labeling requirements of
FIFRA Reg. 1.1471-112

Broadleaf Weeds

Carpeweed

Florida pusley

(Florida purslane)

(Mexican clover)

(Pusley)

Lambsquarters

Pigweeds

(Carelessweed)

(Prostrate pigweed)

(Redroot pigweed)

(Rough pigweed)

(Spiny pigweed)

Purslane, common

Ragweed, common²

Spurge, prostrate¹

Velvetleaf²