Directions for Use Read All Directions Carefully

It is a violation of Federal Law to use this product in a manner inconsistent with the label

Apply Graslan 10P with air equipment calibrated to distribute the pellets uniformly. Apply the pellets at 5 to 10 pounds per acre depending on target brush species.

Grasian 10P may be applied at any time. Treatments become effective after sufficient rainfall has occurred to move the chemical into the root zone where it is taken into the plant. Applications just prior to seasonal rainfall will give the most rapid response. Brush control may continue for several months after application. During this time woody plants may go through repeated defoliations.

Grasian 10P may cause slight injury or suppress desirable grasses. Injury can be minimized by applying when grasses are dormant. If injury to perennial grasses does occur, it will be temporary. Forage grass production usually increases as brush competition is reduced. However, increased grass production is also dependent on adequate rainfall and a sound range management program.

Areas treated with Graslan 10P may be overseeded with a locally adapted variety of tall fescue. Consult your local Range Management Specialist for details on reseeding, such as, varieties, seeding rates, timing and fertilizer programs.

Application Rates:

Grasian 10P is recommended for the control of the following species of undesirable woody plants at the indicated range of application rates.

Higher dosages within the recommended range should be used on deep profile fine textured soils or when treating deep rooted plants.

'Voody Species Controlled		Grasian 10P	
Common Name	Scientific Name	Pounds Per Acre	
Creosotebush	Larrea tridentata	5-10	
On' Sand Shinnery	Quercus havardır	1	

G an **10P** applied as directed will also control several broadleat weeds found in association with undesirable woody plants.

Weeds Controlled		Grasian 10P Pounds Per Acre	
Broomweeds, common perennial Texas	Gulierrezia dracunculoides Xanthocephalem sarothrae Gutierrezia texana	5-10	

Grastan 10P is not recommended for control of persimmon, prickly pear or cholia cactus.

Use Precautions:

Do not apply Grasian 10P on field crops, near desirable trees or shrubs, or areas into which their roots may extend, or in locations where the chemical may be washed in contact with their roots as injury or death may occur.

Do not apply Graslan 10P to newly seeded areas.

Do not apply Grasian 10P under conditions which will cause pellet movement to nontarget areas during application.

Apply Grasian 10P only once per year.

'ian 10P may seriously injure desirable forage legumes such as lespedeza or clover

Lamot cut forage grass for hay from Graslan-treated areas for two years after application.

Do not allow lactating dairy animals to graze on or consume hay harvested from Graslan-treated areas for two years after application

Thoroughly clean all traces of Graslan 10P from application equipment after use. Residues cleaned from application equipment should not be emptied on areas where they will come into contact with the roots of desirable trees, shrubs, plants, or water source.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contact with the leges, or cotting. In case of contact, there with water

ENVIRONMENTAL HAZARD

Keep out of takes, ponds, or streams. Do not contaminate water by clearing of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Dia not contaminate water, food, feed, other pesticides, fertilizer or seeds

Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies and desirable vegetation.

Dispose of empty container in an ir cinerator or landfill approved for pesticide containers, or bury in a safe place away from desirable vegetation.

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

The manufacturer makes no warrant by leaping, our proposition concerning this production its user, when eximple description on the rapps. As statements material correctly with distributing type that discreted

ELANCO

ID 5510



For Control of Brush on Ra in Texas and Oklahoma

Active Ingredient:

Contains 5 pounds active ingredient per 50;

CAUTION: Keep out of reach of panel for additional cautorists.

*Graslan**—the trademark for Elanco Production

1491-115

HEST DOCUMENT AVAILAB

24 PH 5653 A

Elanco Products Company • A Division of it Indianapolis, IN 46206, U.S.A.

EPAF (

Read All Directions Carefully Before Applying Use

roduct in a manner inconsistent with the label

-brated to distribute the pellets uniformly. Apply the pellets at 5 to ush species.

reatments become effective after sufficient rainfall has occurred to $\mathfrak q$ it is taken into the plant. Applications just prior to seasonal rainfall introl may continue for several months after application. During this d defoliations.

suppress desirable grasses. Injury can be minimized by applying arennial grasses does occur, it will be temporary. Forage grass apetition is reduced. However, increased grass production is also d range management program.

verseeded with a locally adapted variety of tall fescue. Consult your etails on reseeding, such as, varieties, seeding rates, timing and

introl of the following species of undesirable woody plants at the

range should be used on deep profile fine textured soils or when

	Grasian 10P		
Vame	Pounds Per Acre		
r tata	5-10		
a rd ii	1		

control several broadleaf weeds found in association with undesir-

	Grasian 10P Pounds Per Acre	
ea dracunculoides ephalem sarothrae ea texana	5-10	

of persimmon, prickly pear or cholla cactus.

mear desirable trees or shrubs, or areas into which their roots may al may be washed in contact with their roots as injury or death

ins which will cause pellet movement to nontarget areas during

- forage legumes such as lespedeza or clover.
- an-treated areas for two years after application.
- Lie on or consume hay harvested from Grasian-treated areas for

P from application equipment after use. Residues cleaned from ed on areas where they will come into contact with the roots of

SUTIONARY STATEMENTS MANS AND DOMESTIC ANIMALS CAUTION

".. eyes or clothing. In case of contact, flush with water

RONMENTAL HAZARD

not contaminate water by cleaning of equipment or disposal

RAGE AND DISPOSAL


r pesticides, fertilizer or seeds

innot be used or chemically reprocessed should be disposed. buried in a safe place away from water supplies and desirable

ator or landfill approved for pesticide containers, or bury in a

ithorities for approved alternative procedures such as limited

or implied, concerning to a product or its use, which is denoted by section reingth approduct applying when users a loncer is



For Control of Brush on Rangeland in Texas and Oklahoma

Active Ingredient: tebuthiuron: N-[5-(1,1-dimethylethyl)-1,3,4thiadiazol-2-yl]-N,N'-dimethylurea 10.0% Inert ingredients: 90.0% Contains 5 pounds active ingredient per 50 pound bag

CAUTION: Keep out of reach of children. See back panel for additional caution statements. *Graslan** — the trademark for Elanco Products tebuthiuron

1491-115

HIST DOCUMENT AVAILABLE

Elanco Products Company • A Division of Eli Lilly and Company Indianapolis, IN 46206, U.S.A.

11 A Rog No. 1471-115