

TREFLAN® E.C. (EPA REG. NO. 1471-35)
TREFLAN® M.T.F.™ (EPA REG. NO. 1471-116)
TREFLAN® 5 (EPA REG. NO. 1471-120)

TREFLAN/SCEPTER® FOR WEED CONTROL IN SOYBEANS
(Do Not Use In California)

TREFLAN/Scepter may be applied as a preplant incorporated tank-mix treatment. Scepter may also be applied as a preemergence overlay treatment or as an early postemergence treatment following a preplant incorporation application of TREFLAN. Use of Scepter in combination with TREFLAN or as a sequential treatment is subject to a number of Special precautions and limitations. Carefully read and understand all special precautions and limitations listed in this supplemental labeling as well as the Scepter label prior to use.

SPECIAL USE PRECAUTIONS

1. Do not use the TREFLAN/Scepter tank-mix or TREFLAN plus overlay of Scepter in the "Northern Use Area" as defined by the Scepter label.
2. The use of Scepter is limited to those states listed on the Scepter label.
3. Certain severe restrictions apply to corn, wheat and other rotational crops following an application of Scepter. Be sure to refer to the Scepter label for complete details of these and other restrictions.
4. A preharvest interval of 90 days is required between the last Scepter application and the soybean harvest.
5. Do not graze or feed treated soybean forage, hay or straw to livestock. Only rotational crops harvested at maturity may be used for food or feed.
6. Do not plant wheat within 4 months following an application of Scepter; or do not plant rice until the spring of the year following a Scepter application.

the spring of the year following a Scepter application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the bootheel of Missouri, Tennessee, North Carolina and states south: field corn may be planted in the spring of the year following a Scepter application unless extreme drought conditions develop and less than 15 inches of rainfall or irrigation is received within 6 months following the date of application.

9. Do not plant cotton or rotational crops other than those listed above within 18 months following an application of Scepter.
10. Crops other than soybeans may be injured by spray, drift or other indirect contact with Scepter.

TREFLAN/Scepter Tank-Mix Directions

DO NOT USE THE TREFLAN/SCEPTER TANK-MIX IN THE "NORTHERN USE AREA" AS DEFINED BY THE SCEPTER LABEL.

Apply TREFLAN/Scepter tank-mix as a preplant incorporated treatment. Follow soil preparation, mixing and application procedures described in the TREFLAN label. Apply TREFLAN/Scepter and shallowly incorporate. Incorporate into soil within 24 hours after application and plant soybeans within 45 days after application. Use equipment that provides uniform incorporation into the top 2 inches of soil. When using a disk, field cultivator, or rolling cultivator, a second pass must be made at an angle to the first pass to ensure thorough incorporation.

If soybeans are planted on beds, apply and incorporate after bed formation.

TREFLAN/Scepter Overlay Directions

DO NOT USE THE TREFLAN/SCEPTER OVERLAY IN THE "NORTHERN USE AREA" AS DEFINED BY THE SCEPTER LABEL.

Apply TREFLAN as a preplant incorporated treatment. Follow soil preparation, application and incorporation procedures described in the TREFLAN label. Apply Scepter after planting but before the crop emerges. Rainfall or overhead irrigation is necessary to move Scepter surface overlay treatments into the weed germination zone. The amount of rainfall or irrigation required depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after a surface applied treatment, a cultivation is recommended to control escaped weeds.

Weeds Controlled (Tank-Mix or Overlay)


TREFLAN/Scepter tank-mix or overlay treatments control the weeds listed on the TREFLAN label for TREFLAN alone (excluding special rate programs) plus these additional weeds:

- | | |
|--|-------------------------|
| Cocklebur, common | Poinsetta, wild |
| Jimsonweed | Prickly sida |
| Mallow, Venice | (Teaweed) |
| Morningglory | Ragweed |
| (Pitted) | (Common) |
| (Smallflower) | (Giant) ¹ |
| Mustard species | Smartweed |
| Nightshade, eastern black ¹ | (Ladysthumb) |
| Pigweed | (Pennsylvania) |
| (Palmer) | Sunflower, common |
| (Smooth) | Velvetleaf ¹ |
| (Tall waterhemp) | |

¹ Eastern black nightshade, giant ragweed, and velvetleaf are controlled by preplant incorporated treatments only.

TREFLAN/Scepter tank-mix or Scepter overlay treatments will aid in the control and reduce competition from these additional weeds:

- Morningglory²
 - (Entireleaf)
 - (Ivyleaf)
 - (Tall)
- Nutsedge, Yellow³
- Panicum, Fall⁴
- Shattercane⁴



4 of 4

Control of these weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of weed germination, depth of weed seed in soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

- ² For best activity on morningglory species, use tank-mix preplant incorporated treatments.
- ³ Use tank-mix preplant incorporated treatments only to aid in control of yellow nutsedge.
- ⁴ TREFLAN alone controls fall panicum and shattercane at increased rates. See the TRFFLAN label for special instructions.

Broadcast Application Rates:

Soil Texture	TREFLAN		Scepter Tank-mix or Overlay (pints)
	E.C./M.T.F. (pints)	or Treflan 5 (pints)	
Coarse	1	0.8	2/3
Medium	1 1/2	1.2	2/3
Fine	2	1.6	2/3

TREFLAN/Scepter Postemergence Directions

Apply TREFLAN as a preplant incorporated treatment. Additional weeds tolerant to TREFLAN may be controlled by using a post emergence application of Scepter. Consult the Scepter product label for application rates additional weeds controlled, application directions and precautions prior to use.

Read the TREFLAN and Scepter labels carefully for additional weeds controlled, directions, precautions, and limitations prior to application.

TREFLAN® E.C., M.T.F.™, 5 (trifluralin, Elanco)
Scepter® (imazaquin, American Cyanamid)

Copyright© 1988 - Elanco Products Company

Elanco Products Company
A Division of Eli Lilly and Company
Indianapolis, Indiana 46285

