

SONALAN/COMMAND®/SCEPTER® COMBINATION FOR WEED CONTROL
IN SOYBEANS
(Do Not Use In California)

Sonalan/Command/Scepter should be applied as a preplant incorporated tank mix. The Sonalan/Command/Scepter combination effectively controls the weeds listed for Sonalan alone for general weed control plus these additional weeds:

Controlled

Common cocklebur	Pigweed (Palmer amaranth)
Common ragweed	Prickly sida (Teaweed)
Eastern Black Nightshade	Velvetleaf (Buttonweed)
Jimsonweed	Venice mallow
Pennsylvania smartweed	

Additional weeds partially controlled by this combination include:

Common sunflower
Morningglory, annual

Control of these weeds may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation.

Application Directions

Tank mixed Sonalan/Command/Scepter should be applied as a preplant incorporated treatment. Apply Sonalan/Command/Scepter up to three (3) weeks prior to planting.

Do not apply aerially or through any chemigation equipment.

Apply Sonalan/Command/Scepter using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10-15 gallons per acre. Use minimum nozzle pressure and boom height while maintaining uniform spray pattern. Coarse sprays are less likely to drift out of the target area than fine sprays.

<u>Soil Texture</u>	<u>Sonalan E.C. (pints)</u>	<u>Command 4 E.C. (pints)</u>	<u>Scepter E.C. (pints)</u>
Coarse	1 1/4 - 2	3/4	1/3
Medium	1 3/4 - 2 1/2	1 1/8	1/3
Fine	2 1/4 - 3	1 1/2	1/3

Precautions

Read the Scepter label carefully for cautions and precautions relating to environmental hazards, planting of rotation crops, sequential program uses of Scepter, harvest restrictions following postemergence treatments of Scepter, use of Scepter in conjunction with Classic®, Canopy®, or Gemini®, grazing restrictions and other directions, precautions and limitations before applying Scepter. The use of Scepter is limited to those states listed on the Scepter label. A postemergence application of Scepter can only be used in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee and Texas following a Sonalan/Command/Scepter preplant incorporated treatment.

Special Precaution

Off-site movement of spray drift or vapors of Command herbicide can cause foliage whitening or yellowing of some plants. Prior to making application, read and strictly follow all precautions and application instructions on the Command label. Read the Command label carefully for additional directions, cautions and limitations prior to use.

Read the Sonalan label carefully for additional directions, precautions and limitations before applying these products.

161****/EC**36/LABELS/US/****/303

Sonalan® (ethalfluralin, Elanco)
 Canopy® (chlorimuron ethyl plus metribuzin, DuPont)
 Classic® (chlorimuron ethyl, DuPont)
 Command® (FMC)
 Gemini® (chlorimuron ethyl plus linuron, DuPont)
 Scepter® (imazaquin, American Cyanamid)

Elanco Products Company
 A Division of Eli Lilly and Company
 Indianapolis, Indiana 46285