Directions for Use

See attached literature for complete directions for use.

Read all directions carefully before applying.

It is a violation of Federal Law to use this product in a manner inconsistent with às labeling.

Precautionary Statements

Hazards to Humans and Domestic Animals WARNING

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid inhaling dust from product. In case of contact, flush with water.

Environmental Hazards

Do not contaminate any body of water, ponds or streams as death or injury may occur to vegetation irrigated by such. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Storage: The herbicidal properties of Spike Dry Flowable require caution in handling, storage and transportation of this product. Store in original container only. In case of leak or spill, contain material and discose as waste. Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal

Container Disposal: Triple rinse (or equivalent). Do not reuse. Then puncture and dispose of in a sanitary landfill, away from desirable plants, trees and water supply, or by Incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke,

The manufacturer makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed. Spike Dry Flowable must be applied according to Elanco's written instructions, including, but not limited to, recommended rates. Failure to do so may result in poor weed control or plant injury. Elanco expressly disclaims any warranty, either express or implied, for the use of Sorke Dry Flowable, alone or in combination with other products, when that use is not in strict compliance with Elanco's written

เข็บ ปี340 ใหม่

FN 3084

Indianapolle, IN 46285, U.S.A.

Elanco Products Company . A Division of Eli Lilly and Company

ACCEPTED

APR 1 5 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the posticide realstored under EPA Reg. No.//

ELANCO

ID 5943



TM

Dry Flowable

Net Weight 4 Pounds

A preemergence and postemergence herbicide for control of brush and weeds in such areas as:

- Around Industrial buildings
- · Ditchbanks · Railway roadbeds & ballast

Lumberyarda

- Firebreaks
- · Raliroad rights-of-way

Tank farms

- Fence rows
- Railroad yards

- Airport runways
- . Utility substations and rights-of-way
- · Road shoulders where no vegetation is desired
- At the base of transmission towers and poles
- At the base of highway guardralis, signposts, and markers
 Under asphalt and concrete pavements where no future iandscaping is planned

The degree and duration of control may vary with the amount of chemical applied, soil texture, and other conditions.

Active Ingredient:

lebulhluron* M5-(1,1-dimethylethyl)-

Contains 3.4 pounds active ingredient per 4 pound bottle.

*Spike"-the registered trademark for Elanco Products tebuthkiron

Keep Out of Reach of Children.

WAHNING

See side panel for additional precautionary statements.

EPA Reg. No. 1471-147

EPA Est. 1471-IN-5

Bry Flowable

ELANCO

A preemergence and postemergence herbicide for control of brush and weeds

Active Ingredient:
tebuthiuron* M5-{1,1-dimethylethyl}-1,3,4thiadiazol-2-yl|-N,N-dimethylurea 85.0%
Inert Ingredients 15.0%
Contains 3.4 pounds active ingredient per 4 pound bottle

*Spike®—the registered trademark for Etanco Products
tebuthiuron.

EPA Reg. No. 1471-147 Elanco Products Company A Division of Eli Lilly and Company Indianapolis, IN 46285, U.S.A.

Copyright[©] 1984 by Elanco Products Company

Literature Issued November 13, 1984 PI 2950 AMP




Table of Contents Page Precautionary Statements 2 Storage and Disposal 3 Precautions 4 Directions for Use 6 Total Vegetation Control 6 Application Directions 10 Woody Plant Control 12 Broadcast Application 17 Banded Application 18 Individual (Spot) Application 20

The manufacturer makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed. Spike Dry Flowable must be applied according to Elanco's written instructions, including, but not limited to, recommended rates. Failure to do so may result ir. poor weed control or plant injury. Elanco expressly disclaims any warranty, either express or implied, for the use of Spike Dry Flowable, atone or in combination with other products, when that use is not in strict compliance with Elanco's written recommendations.

Precautionary Statements

Keep Out of Reach of Children.

Hazards to Humans and Domestic Animals WARNING

Harmful II swallowed. Avoid contact with skin, eyes or

clothing. Avoid inhaling dust from product. In case of contact, flush with water.

Environmental Hazards

Do not contaminate any body of water, ponds or streams as death or injury may occur to vegetation irrigated by such. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Storage: The herbicidal properties of Spike Dry Flowable require caution in handling, storage and transportation of this product. Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Do not reuse. Then puncture and dispose of in a sanitary landfill, away from desirable plants, trees and water supply, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

3

30

 \frown

.

Precautions



Spike Dry Flowable is intended for noncropland vegetation control. It is an extremely active herbicide which will kill trees, shrubs and other forms of desirable vegetation having roots extending into

the treated area. Feeder roots of many species of desirable vegetation extend many feet beyond the drip line of the branches, and a very small amount of Spike Dry Flowable in contact with one feeder root of a tree, shrub or other desirable vegetation may cause serious injury or death to the entire plant.

An arboriculturist (tree expert) should be consuited to help you to determine if the area of proposed application is free of all roots of desirable vegetation. The effect of Spike Dry Flowable on desirable vegetation may be irreversible and its presence in the soil may prevent growth of other desirable vegetation for some years after application.

Read the entire label before using Spike Dry Flowable to determine if this product is suitable for the desired purpose.

Do not use Spike Dry Flowable on areas such as walks, driveways, streets, lawns, patios, tennis courts, swimming pools, cemeteries, or other landscaped areas, or under asphalt or concrete pavement where future landscaping is planned. Do not apply on field crops. Do not apply on any area into which the roots of field crops or other desirable vegetation may extend. Roots of trees, shrubs, and other desirable vegetation

may extend far beyond the drip line of the plant's branches.

Avoid nontarget drift or product movement. Do not apply when winds are gusty or under any other condition which will allow drift or product movement. Do not apply to areas where soil movement by water erosion and/or natural or mechanical means is likely. Do not apply to areas where wind is likely to cause soil movement unless a soil sealant is used. Drift or any form of product movement from treated areas may cause damage to any vegetation to which treatment is not intended.

Ditchbank Usage—Do not apply Spike Dry Flowable to any portion of the ditchbank that will come into direct contact with water as movement of Spike Dry Flowable in this water to non-target plant species may result in the injury or death of those plants. Do not apply on ditches used to transport irrigation or potable water. Keep from contact with other pesticides and seeds.

Thoroughly clean all traces of Spike Dry Flowable from application equipment after use. Do not empty residues cleaned from application equipment on areas where they may come in contact with the roots of desirable vegetation or the water source for such vegetation.

Woody Plant Control—Grazing is allowed in areas receiving band or individual plant treatments with 4.70 pounds per acre or tess of Spike Dry Flowable. In areas receiving band or individual plant treatments with 4.70 pounds per acre or less of Spike Dry Flowable, grass may be cut for hay one year after application.

á

Spike Dry Flowable may injure or suppress certain herbaceous vegetation in the treated area. Therefore, do not apply where such injury cannot be tolerated. Do not apply broadcast applications of Spike Dry Flowable where torage or maintenance of a grass cover is desired. Injury to most herbaceous perennials is reduced if Spike Dry Flowable is applied when this vegetation is dormant.

Directions for Use

Read all directions carefully before applying. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Total Vegetation Control

Spike Dry Flowable is a preemergence and postemergence herbicide for total control of vegetation in such noncropland areas as: airport runways, utility substations and rights-of-way, and concrete pavements where no future landscaping is planned, at the base of highway guardrails, sign posts and markers, at the base of transmission towers and poles, around industrial buildings, lumberyards, railroad yards, ditchbanks, firebreaks, and fence rows.

For total vegetation control in areas not treated the previous season with Spike Dry Flowable or other residual herbicides, apply Spike Dry Flowable prior to or just after emergence of plants as follows:

At 4.75 pounds per acre, Spike Dry Flowable will control the following:

Alfalfa

Aster, heath

Aster, while heath Barley, little

Bedstraw Bluegrass, annual Bluegrass, Kentucky Bouncingbet Bromegrass, downy Bromegrass, ripgul Bromegrass, smooth Broomsedge Buffelgrass Burclover Buttercup. smallflower Camphorweed Carrot, wild Catsear, spotted Cheat Chickweed Clover, red Cocklebur Creeper, Virginia Crowlootgrass Dock, curly Doglennel Fescue Fescue, rattail Fiddleneck, coast Filaree Filaree, redstem Fleabane, annual Foxtail Gaillardia. rosering Gerantum. Carolina Goldenrod

Grape Gumweed Hemlock, poison Henbit Honeysuckle. Japanese Horseweed Knapweed Kochla Lambsquarters Lupine Medic, black Morningglory Mullein, common Nightshade. silverleas Oat, wild Panicum, Texas Pepperweed. Virginia Pigweed Plantain, buckhorn Puncturevine Ragweed, glant Raspberry, red Ryegrass, Italian Sedge, annual Shepherdspurse Sida, prickly Sowthistle, annual Spikeweed Spurge Spurge, spotted Starthistle. yellow

Strawberry Sunflower, common Telegraphplant Timothy

Trumpetcreeper Velvelgrass Vetch Witchgrass

At 7.0 pounds per acre, Spike Dry Flowable will control the following:

Awngrass, triple Barley, foxtail Bromegrass, Japanese Canarygrass, reed Carpetweed Chicory Cinquefoil, common

lvy, poison Reed, common Sandbur, field Smartweed, swamp Sowthistie. perennial Spurge, prostrate Sumac, staghorn Sweetclover, white

At 9.5 pounds per acre. Spike Dry Flowable will control the following:

Barnyardgrass Craborass

Oxtongue, bristly Poorjoe

(Ichorass Johnsongrass, seedling Lovegrass

Purslane, common Ragweed, common Saltbush

Orchardgrass

Sedge, bull At 18.75 pounds per acre, Spike Dry Flowable will con-

troi the following: Bermudagrass

Lookingglass, Venus Vaseygrass

Cudweed Dallisgrass

For the maintenance of total vegetation control in noncropland areas east of the Rocky Mountains which were treated the previous season with Spike Dry Flowable or other residual herblcides, apply Spike

Dry Flowable prior to or just after emergence of plants as follows: (Some of the species listed may show erratic control depending on the time between application and weed germination.)

At 2.0 pounds per acre, Spike Dry Flowable will control the following:

Bluegrass, annual Bluegrass, Kentucky Carrot, wild

Parsnip, wild Pepperweed Pigweed

Chickweed, common Croton

Ragweed, common Smartweed, Pennsylvania

Fleabane, annual Horseweed Mullein Panicum, fall

Sweetclover Thistle, Canada Woodsorrel, yellow

At 3 pounds per acre. Spike Dry Flowable will control the following:

Goldenrod

Spurge

In areas of rainfall greater than 25 inches per year. the 3 pounds per acre maintenance rate should be used for all weed species listed above.


For the maintenance of total vegetation control in noncropland areas west of the Rocky Mountains which were treated the previous season with Spike Dry Flowable or other residual herbicides, apply Spike Dry Flowable prior to or just after emergence of plants as follows: (Some of the species listed may show erratic control depending on the time between application and weed germination.)


At 1.5 pounds per acre, Spike Dry Flowable will control the following:

Bassia, fivehook Cheat

Cudweed Foxtail

9





Lettuce, prickly

Oat, wild Oxtongue, bristly Pigweed

Plantain

At 2.0 pounds per acre, Spike Dry Flowable will control the following:

Buttercup

Canarygrass, reed Knapweed, Russian

Knotweed Mallow Mustard Ragweed, western Starthistle.

yellow Telegraphplant

Ryegrass, annual

Shepherdspurse

Saitbush

Witchgrass

At 3.0 pounds per acre, Spike Dry Flowable will control the following:

Barley Gumweed Sida, alkali Smartweed, swamp

Puncturevine

In areas of rainfall greater than 25 inches per year, the 3 pounds per acre maintenance rate should be used for all weed species listed above.

Application Directions

Apply Spike Dry Flowable in 15 to 150 gallons of water per acre before or during the period of active growth of plants to be controlled. Initial control is enhanced by rainfall.

In areas of low annual rainfall (less than 15 inches per year) Spike Dry Flowable should be applied prior to the time of year when the predominant portion of that rainfall occurs. A minimum of 1 to 1 1/2 inches of rainfall is required to activate Spike Dry Flowable and place it in the primary weed seed germination zone.

To maximize performance under a variety of conditions, Spike Dry Flowable is recommended for use in tank-mix combination with any of the following products: amitrole, Atratol® 80W, Banvel® 720, Banvel® W.S., Karmex® 80W, MSMA, paraquat, Princep® 80W. Roundup®, Surflan® 75W, 2.4-D or Oust®. Where applications are made to existing vegetation, the contact or burning properties of paraquat or MSMA provide the rapid top kill while Spike Dry Flowable gives the residual long-term control desired. Applications to areas which are infested with certain difficult to control perennial weeds (such as johnsongrass, bermudagrass, quackgrass, horsetail, bindweed, dandelion or nutsedge) will benefit from tank-mixing Spike Dry Flowable with Roundup, amitrole, Banvel, Banvel 720, or 2,4-D. The addition of Surflan, Oust, Karmex, Atratol or Princep will improve Spike's performance on certain annual broadleaf and grass weeds such as foxiall, kochia, Russian thistle, or seedling Johnsongrass.

Read the Spike Dry Flowable label and labels of products to be tank-mixed carefully before using. Note all warnings, cautions, precautions, and limitations of warranty on all labels.

Apply with any sprayer that will apply the spray uniformly. Check the sprayer before and during use to insure proper calibration and uniform application.

Surflan®-oryzafin, Elanco Products Company
Abatol®-atrazine and prometon, Ciba-Geigy
Banvat®-dicamba, Velsicol Chemical Corporation
Karmex®-dicron, E. I. DuPont De Nemours & Company
Princep®-almazine, Ciba-Geigy
Roundup®-glyphosate, Monsanto Company
Oust®-suttometuran methyl, E. I. DuPont De Nemours & Company

10

To mix, fill spray tank half full of water. Start agitation and continue during the entire mixing process. Add required amount of Spike Dry Flowable and allow to mix when tank mixing. If additional product is a wettable powder, add to tank and allow to mix thoroughly. If additional product is a liquid, add slowly while filling remainder of tank with water. Continuous agitation in the spray tank is required to keep the materials in suspension throughout application.

Agitate by mechanical or bypass (hydraulic) means in the spray tank. If bypass or return agitation is used, it should terminate at the bottom of the tank to minimize loaming.

For treating small areas, a tank type hand sprayer or sprinkling can may be used. Before application determine the amount of water and chemical necessary to cover uniformly the area to be treated. Shake or stir frequently.

Woody Plant Control

Spike Dry Flowable is an effective herbicide for the control of brush and vines. Spike Dry Flowable can be applied either as a broadcast spray, banded application or as an individual plant treatment depending upon the size, density and location of brush to be controlled.

Spike is to be applied to the soil (Not the Foliage!) where it is absorbed by the roots of plants. Effects are slow to appear and will not become apparent until sufficient moisture has carried Spike Dry Flowable into

the root zone. The time required to achieve control is dependent on soil type, amount of rainfall and rooting depth of target species. Some species may go through several defoliations and refoliations over a period of approximately two to three years prior to dying.

Spike Dry Flowable can be applied anytime except when the ground is frozen or the soil is saturated with moisture. For optimum results, applications should be made just prior to the resumption of active seasonal growth in the spring and/or periods of rainfall. For applications made in the late summer or early fall in areas of average annual rainfall of greater than 25 inches, higher rates should be used and inconsistent control may result on densely infested brush areas and hard to control species.

Spike Dry Flowable may be used on cut brush but for optimum results time should be allowed for the brush to resprout to a height of approximately 5 feet prior to application. Spike requires an actively growing plant to be effective. The larger the resprouts the more Spike that will be taken up by the plant and the more effective and consistent the control will be.

For the control of woody plants and vines, the following rates of Spike Dry Flowable are recommended. These rates can vary depending upon soil type, rainfall, time of application and size/density of the woody plants.

Spike Dry Flowable applied at the rate of 1.25 pounds per acre will control the following species.

Haplopappus tenuisectus Burroweed Larrea tridentata Creosotebush

12



Mimosa biuncitera Wait-a-minute-bush Spike Dry Flowable applied at the rate of 2.50 pounds per acre will control the following species.

Ailanthus altissima Aloysia lycioides Artemisia tridentata Carva glabra Celtis occidentalis Datura discolor Lycium berlandieri

Tree-of-tieaven Whitebrush Big sagebrush Pianut hickory Western hackberry Desert thornapple Berlandier wolfberry Red mulberry

Morus rubra Pínus monticola Pinus spp. Prunus emarginata Rhus alabra Robinia pseudoacacia Rosa multillora Salvia leucophylla Symphoricarpos orbiculatus

Western white pine 9ni9 Bitter cherry Smooth sumac Black locust Multiflora rose Black sage

Buckbrush Ulmus americana American elm Vaccinium spp. Blueberry Gaylussacia spp. Huckleberry

Spike Dry Flowable applied at the rate of 3.50 pounds per acre will control the following species.

Abies balsamea Acacia farnesiana Acer saccharum Alnus rugosa Belula populifolia Carya taxana Celtis pallida Condalia obtusifolia

Balsam tir Huisache Sugar maple Speckled alder Gray birch Black hickory Granieno Lotebush condalia llex vomitoria Larix laricina Picea glauca Populus balsamilera Populus deltoides Quercus douglasii Quercus marilandica Quercus stellata Salix spp. Schaelleria cuneilolia Spiraea tomentosa Úlmus alata

Yaupon Tamarack White spruce Balsam poplar Eastern cottonwood Blue Oak Blacklack oak Post oak Willow Desert yaupon Hardhack Winged elm

Spike Dry Flowable applied at the rate of 4.70 pounds per acre will control the following species.

Guaiillo

Catclaw acacia

Acacia berlandieri Acacia greggii Acacia rigidula Acacia tortuosa Acer negundo Adenosioma fasciculatum Alnus rubra Campsis radicans Carva ovata Cercocarous betuloides Colubrina texensis Condalia obovata Cornus drummandii Crataegus spp. Evsenhardtia texana Fagus grandifolia

Jalropha dioica

Leucophyllum

Irulescens

Blackbrush acacia Twisted acacia Boxelder Chamise Red alder Trumpelcreeper Shaobark hickory Birchleaf mountain-mahogany Texas colubrina Bluewood condalla Roughleaf dogwood Hawlhorn Texas kidneywood American beech

Leatherstem Cenizo

(Texas silverleaf)

14

Liquidambar styraciflua Parthenocissus auinauelolia

Populus grandidentala Porlieria angustifolia Prunus virginiana Pseudotsuga menziesii

Pueraria lobata Quercus dumosa

Quercus palustris Quercus rubra Quercus virginiana Rhus Ivohina Rubus allegheniensis

Salvia ballotaeflora

Sweetgum

Virginia creeper Bigtooth aspen

Guayacan Common chokecnerry

Douglas fir Kudzo California scrub oak Pin oak Red oak Live oak

Staghorn sumac Allegheny

Bigleat maple

b' ckberry Shrubby blue salvia

Spike Dry Flowable applied at the rate of 6.0 pounds per acre will control the following species.

Acer macrophyllum Acer platanoides Acer saccharinum Baccharis spp. Cornus florida Fraxinus pennsylvanica Gaultheria shallon Juniperus virginiana Lantana camara Liriodendron tulipilera Melaleuca quinquenervia Pinus banksiana Pinus echinata Pinus resinosa Pinus virginiana Platanus occidentalis

Norway maple Silver maple Groundsel tree Flowering dogwood Green ash Salal Eastern redcedar Lantana Tuliptree Melaleuca Jack pine Shortleaf pine Red pine Virginia pine American sycamore

Prunus serolina Black Cherry Quercus alba White cak Rubus laciniatus Evergreen blackberry Rubus occidentalis Black raspberry Schinus

Brazilian terebinthifolius peopertree

Spike Dry Flowable applied at the rate of 7.0 pounds per acre will control the following species.

Acer circinatum Arciosiaphylos patula Ceanothus cuneatus Ceanothus leucodermis

Cartaegus crus-galli Elagagnus angustitotia Fraxinus americana

Ligustrum spp. Rhus laurina Sapium sebilerum Smilax rotundifolia

Ulmus parvilolia Ulmus rubra

Vine maple Greenleaf manzanita

Wedgeleaf ceanothus Whitelhorn

chaparral Cockspur hawthorn Russianolive White ash Privet Laurel sumac Taliowtree Common greenbrier

Chinese elm Slipper, elm

Broadcast Application

Apply Spike Dry Flowable in 15 to 150 gallons of water per acre with any properly calibrated herbicide sprayer. Check the sprayer before and during use to insure proper celibration and uniform application. Add the recommended amount of Spike Dry Flowable to clean water in the spray tank during the filling operation. Material must be kept in suspension at all times by constant agitation. Agitate by mechanical or bypass (hydraulic) means in the spray tank. If bypass or return

agitation is used, it should terminate at the bottom of the lank to minimize foaming.

For treating small areas, a tank type hand sprayer may be used. Before application, determine the amount of water and chamical necessary to cover uniformly the area to be troated. Shake or stir frequently.

Do not apply broadcast applications of Spike Dry Flowable where forage or maintenance of a grass cover is desired.

Banded Application

Spike Dry Flowable is recommended for the control of woody plant species in noncropland areas (such as utility, railroad, and pipeline rights-of-way, ditchbanks and fence rows) by application of a series of parallel bands to the soil surface. Individual bands should be spaced at intervals from 4 to 10 feet and at the currently labeled rate range of 2.5 to 7.0 pounds per acre depending on the woody species to be controlled. Actual herbicide bands should be kept as narrow as possible during application to achieve minimal injury or control of herbaceous vegetation. Apply Spike Dry Flowable to the soil surface in 5 to 75 gollons of water per acre in a series of parallel bands with spacing between bands ranging from 4 to 10 feet. In areas such as brush infested fence rows on utility rights-of-way, a single band may be applied. Control is dependent upon root systems intercepting bands. Therefore larger stems should be treated individually when using single bands.

Band spacing should be selected based on the size

of the woody plants in the area to be treated and the amount of injury or control of herbaceous vegetation that can be tolerated.

Where control of young or seedling woody plants is desired, bands should be spaced closer together. This will achieve maximum exposure to their limited root systems. Where larger more mature woody species are to be controlled, bands should be spaced at the wider end of the recommended spacing range.

In addition to allowing adequate exposure of the more extensive root systems of these larger woody species for control, use of the wider spacings will further reduce injury or control of herbaceous vegetation in the area of treatment.

Within the treated band nearly all vegetation, woody and herbaceous, will be killed. Some herbaceous vegetation close to the treated band with roots extending into it may be severely injured or killed. However, since herbaceous species tend to have restricted root systems, most species outside the treated band will not be affected. Banded applications in areas of steep terrain should be applied across existing stopes in order to prevent soil erosion.

Apply with equipment designed to deliver the spray uniformly in the bands. To maintain the integrity of the individual herbicide bands, straight stream nozzles fitted with internal stabilizing vanes or their equivalent are recommended. Operating pressures should also be kept as low as will provide uniform delivery of the spray solution. Pressures in the range of 10 to 40 psi should be adequate. Pressures in excess of 40 psi will tend to cause the individual bands to break up.

19

9

When applications are made in an area where nozzles are elevated above the soil surface more than 5 feet, breakup of the individual spray streams may occur. If conditions do not permit delivery of intact spray streams to the soil surface, this method of application should not be used.

Fill the spray tank half-full of water. Start agitation and continue during entire mixing and spraying operation. Add the required amount of Spike Dry Flowable and allow it to mix thoroughly white completing the spray tank filling. If hand held or back pack type sprayers are used, shake vigorously after filling and periodically during application to maintain product suspension. A master shut-off switch for the entire spraying system and nozzle check valves are recommended on commercial spray equipment.

Material must be kept in suspension at all times by continuous agitation. Agitate by mechanical or bypass (hydraulic) means in the spray tank. If bypass or return agitation is used, it should terminate at the bottom of the tank to minimize foaming. Check the sprayer frequently before and during use to insure proper calibration and uniform application.

Individual (Spot) Application

Spike Dry Flowable may be applied in high or low volumes of water to selectively control individual woody plants. Recommended rates will vary depending upon site conditions with the higher rates needed for difficult to control species, large plants, heavier soils, fall applications and cut brush. Consult your local Elanco Spike distributor to determine the best rates for your location.

For high volume applications, mix 1 pound of Spike Dry Flowable in enough water to make 10 gallons of solution. Apply 10 ounces of material to the soil per every 2 to 4 inches of stem diameter.

For low volume applications, mix 1 pound of Spike Dry Flowable in enough water to make 1 gallon of solution. Apply 1 ounce of material to the soil per every 2 to 4 inches of stem diameter.

When treating large stems, apply the multiple treatments in even specing around the stem.

For applying Spike Dry Flowable in banded or individual plant treatment, two pieces of equipment are suggested; the Solo Model 425 back pack sprayer for both banding and individual plant treatment and the Spot Gun for individual plant treatment.

The Solo sprayer is prepared for spraying by adding the pre-slurried contents of a 4 pound bag of Spike Dry Flowable and water to the tank. Fill to capacity with additional water and shake vigorously. Equip the Solo sprayer with a 0003-SS straight stream nozzle and the Solo pressure regulator with the green (10 psi) pressure limiting spring. To band Spike Dry Flowable at 5 pounds per acre, walk at 3 mph (264 feet per minute) with the Solo on continuously and space the bands 5 feet apart. Adjust the rate and walking speed according to the brush species and conditions encountered. For individual plant treatment with the Solo, apply a 1.5 second shot for every 1 to 2 inches of stem diameter at the base of unwanted woody plants.

The Spot Gun is prepared for individual plant treatment by mixing 2 pounds of Spike Dry Flowable in

sufficient water to obtain 1 gallon of spray solution. Set the Spot Gun to deliver 8 milliliters of this solution for every 1 to 2 inches of stem diameter at the base of the unwanted woody plants. For application on steep slopes or other sensitive areas, the Spot Gun can be equipped with a soil probe to allow injection of the Spike Dry Flowable solution beneath the soil surface. Placement at a soil depth of 2 to 4 inches will eliminate any surface movement and reduce injury to herbaceous vegetation.

At the prescribed rates, a 4 pound bag of Spike Dry Flowable will treat approximately 950 stems 1 to 2 inches in diameter. Because of its non-volatile nature and low potential for drift this Spike application technique can be used for treating unwanted woody plants growing on non-cropland areas adjacent to sensitive crops. (See label precautions.) A white spot should be visible at the base of each treated stem which should aid in inspection of the completed work.

Precaution: Do not use Spike Dry Flowable in this manner in any area where desirable species are in the vicinity of the plants to be eliminated. A small amount of Spike Dry Flowable in contact with the roots of desirable trees or other woody species may cause severe injury or death. The roots of such plants may extend far beyond their drip lines.

Spike Dry Flowable will injure or control other herbaceous vegetation in the treated area. Therefore, do not apply where such injury cannot be tolerated. See the list of herbaceous vegetation controlled by Spike Dry Flowable under the Total Vegetation Control section of this booklet.

P23 BLANK