


10F4
1125

1471-168

(11/14/86)



ELANCO

TURF FERTILIZER

contains

TEAM™ 1.25%

A GRANULAR FERTILIZER CONTAINING TEAM™ FOR THE PREEMERGENCE CONTROL OF CRABGRASS, FOXTAIL, GOOSEGRASS, POA ANNUA AND OTHER ANNUAL WEED GRASSES IN ESTABLISHED TURFGRASSES.

Active Ingredients:

N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine.....0.82%
 trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-
 p-toluidine.....0.43%

Inert Ingredients.....98.75%

Contains 0.625 pounds active ingredient per 50 pound bag.

TEAM™ (benefin plus trifluralin, Elanco)

GUARANTEED ANALYSIS

TOTAL NITROGEN(N).....24.0%

AVAILABLE PHOSPHORIC ACID(P₂O₅)..... 6.0%

SOLUBLE POTASH(K₂O).....12.0%

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash

1052*C*/G*1.25/LABELS/US/****/1

thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Statements of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

If swallowed: Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention.

Environmental Hazards

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE - Read All Directions Carefully Before Applying

Elanco's Turf Fertilizer containing TEAM can be used on the following established cool and warm season turfgrasses:

Established Cool Season Turfgrasses

- Bentgrass Fairways
- Kentucky bluegrass
- Perennial ryegrass
- Fescues

Established Warm Season Turfgrasses

- Bahagrass
- Bermudagrass
- Centipede grass
- St. Augustine grass
- Tall Fescue
- Zoysiagrass

Elanco fertilizer containing TEAM is a preemergence herbicide which is applied to established turfgrasses for the control of the winter annual grass: annual bluegrass (Poa annua); and summer annual grasses: smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfootgrass), barnyardgrass (watergrass) and green and yellow , foxtail.

This product should be applied one to two weeks prior to the anticipated germination of the target annual grass weeds. Apply using a properly calibrated granular fertilizer spreader which will uniformly distribute the product over the target area. Avoid skips and excessive overlaps as poor control or turfgrass injury may occur. This product will not harm most nearby ornamentals.

Rates and Frequency of Application

| | <u>Cool Season Turfgrasses</u> | <u>Warm Season Turfgrasses</u> |
|---------------------------|--|--|
| Summer Annual Grass Weeds | 160-240 pounds/acre (3.7 - 5 1/2 pounds/1000 sq. ft.) | 240 pounds/acre (5 1/2 pounds/1000 sq. ft.) |
| Winter Annual Grass Weeds | 160 pounds/acre | 240 pounds/acre |


In areas with extended growing seasons or where desired, a second application at the recommended rate can be made 8 to 10 weeks following the initial application on cool season turfgrasses and 10 to 12 weeks following the initial application on warm season turfgrasses.

Do not reseed treated areas for a minimum of 8 weeks after an application of 160 pounds per acre or 12 to 16 weeks after an application of 240 pounds per acre.

Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

General Use Precaution

Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipede grass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application.



DISCLAIMER OF WARRANTIES

The manufacturer makes no warranties concerning this product or its use, which extend beyond the description on the label. All other warranties, express or implied, including but not limited to any implied warranties of merchantability and fitness for a particular purpose, are disclaimed. All statements concerning this product apply only when used as directed.

LIMITATION OF DAMAGES


Elanco's liability, whether in contract, warranty, tort, negligence, strict liability or otherwise, should not exceed the return of the amount of the purchase price of the turf fertilizer containing TEAM and under no circumstances should Elanco be liable for special, indirect or consequential damages.

TEAM™ (benefin plus trifluralin, Elanco)

EPA Est. No. XXXXX-XX-XX
EPA Reg. No. 0000-000

Elanco Products Company
A Division of Eli Lilly and Company
Indianapolis, IN 46285, U.S.A.

Team Turf Fertilizer/1.25
11/14/86



1052*G*/G*1.25/LABELS/US/****/4