

Kills Fleas, Ticks, Lice

Contents

Active ingredients:

- Malathion.....0.5%
[O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate]
- Pyrethrins.....0.1%
- Technical Piperonyl Butoxide.....0.4%
[equivalent to 0.32% (butylcarbityl) (6-propylpiperonyl) ether and 0.08% of related compounds]
- 2,2'-Methylenebis-(4-chlorophenol).....0.25%
[dichlorophene]
- Isopropanol.....1.9%
- Petroleum distillate.....18.1%

Inert ingredients.....75.75%

CAUTION: Avoid breathing mist. Avoid contamination of foodstuffs. Wash hands after applying. Do not expose container to temperatures in excess of 130° F. or to extreme cold. USDA Reg. No. 1516-47

FLEACIDE® - M

SPRAY



May be used on Dogs and Cats

CAUTION: Keep out of reach of children. See left panel for additional cautions.

Manufactured by

CURTS

Laboratories, Inc.

KANSAS CITY,
MISSOURI, U.S.A.
64105

SOLD ONLY TO VETERINARIANS

Checks Itching — Checks Odor

Directions

Hold can upright 2 to 4 inches from the dog or cat. Point nozzle in direction spray is desired. Release spray by depressing valve. Direct spray over the animal until the entire coat is damp. A 30 to 45 second application is usually sufficient to treat an average size dog; for cats, 10 seconds. On heavily coated animals, ruffle the coat and spray against the lay of the hair. For best results in controlling ticks, spray directly on the tick. Thoroughly spray interior of kennel or other sleeping quarters. Repeat treatments should be limited to not more often than once weekly. Nervous animals soon become accustomed to the hissing sound and cool sensation of this spray.

NET WT.: 12 OUNCES

