

PHOENIX BRAND

METHYL PARATHION* EC-46

EMULSIFIABLE

ANTIDOTE STATEMENT

DANGER

Poisonous if swallowed, inhaled, or absorbed through the skin. Rapidly absorbed through the skin! Repeated inhalation or skin contact may progressively increase susceptibility to methyl parathion poisoning without symptoms. If you suspect exposure, see a doctor. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wear a mask or respirator of a type passed by the U.S. Dept. of Agriculture for methyl parathion protection. Wear natural rubber gloves, goggles, and clean protective clothing daily. Destroy gloves and replace frequently. Do not contaminate feed or foodstuffs. Not for use or storage in or around the home.

FIRST AID

EXTERNAL: For skin contact, immediately remove contaminated clothing and wash thoroughly with soap and water. For eyes, flush immediately with water for at least 10 minutes.

INTERNAL: CALL A PHYSICIAN. Induce vomiting by giving warm salt or soap water until vomit fluid is clear. Have victim lie down and keep quiet.

NOTE TO PHYSICIANS: Atropine and/or 2-PAM are antidotal. Methyl parathion is a cholinesterase inhibitor. Warning symptoms include weakness, headache, pin-point pupils, salivation, nausea, vomiting, diarrhea and abdominal cramps.

DANGER

This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed. Apply this product only as specified on this label.

Do not contaminate lakes, streams or ponds with this material.

TREATED FIELDS MUST BE POSTED.



U.S.D.A. Reg. No. 1526-194

ACTIVE INGREDIENTS:	
0,0-dimethyl 0-p-nitrophenyl thiophosphate*	46.00%
Aromatic Petroleum Solvent	48.00%
INERT INGREDIENTS:	6.00%
	100.00%

*Methyl parathion

CONTAINS 4 POUNDS METHYL PARATHION PER GALLON AT 60°F

FLAMMABLE MIXTURE

Keep away from heat or open flame. Read entire label before use and comply with all precautions and directions.

DANGER

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Prevent drifting to crops intended as feed for dairy or slaughter animals.

Vacated areas should not be reentered until the drifting insecticide and volatile residues have dissipated.



DANGER POISON



DANGER: KEEP OUT OF REACH OF CHILDREN.

SEE WARNING AND ANTIDOTE STATEMENTS ON SIDE PANEL



Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

DIRECTIONS FOR USE

For ground application dilute in 10 to 30 gallons of water per acre. For air application dilute in 30 to 10 gallons of water per acre

CROP	INSECT PESTS	MAXIMUM RATE	LIMITATIONS
Alfalfa	Aphids	1½ quarts per acre	Do not apply within 20 days of harvest
	Armyworms		
	Leafhoppers	-or-	Do not apply within 15 days of harvest
	Mites	1 quart per acre (in California use 3/4 pint per acre)	
Barley, wheat	Aphids	1/2 pint per acre	No time limitation
	Armyworms	-or-	
	Leafhoppers	1½ pints per acre	Do not apply within 15 days of harvest
		-or-	
		1½ quarts per acre	Do not apply after heads begin to form
Broccoli	Aphids	1 pint per acre	Do not apply within 7 days;
Brussels sprouts			10 days for cabbage
Cabbage	Aphids	1½ quarts per acre	Do not apply within 21 days of harvest
Cauliflower	Cabbage looper		
Carrots	Aphids	1 quart per acre	Do not apply within 15 days of harvest. Do not use treated tops for food or feed.
	Armyworms		
	Flea beetles		
	Leafhoppers		

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CROP	INSECT PEST	RATE	LIMITATIONS
Cotton	Aphids	3 quarts per acre	Do not apply within 5 days of handpicking. (Workers entering fields within 24 hrs. should wear protective clothing.)
	Fleahoppers		
	Lygus bugs		
	Thrips		
	Cotton leaf perforator		
Lettuce	Aphids	1 quart per acre	Do not apply within 21 days of harvest
	Armyworms		
	Leafhoppers		
Cabbage worms			
Onions	Thrips	7/8 quart per acre	Do not apply within 15 days of harvest
Potatoes	Aphids	1½ quarts per acre	Do not apply within 5 days of harvest
	Armyworms		
	Flea beetles		
	Leafhoppers		
Safflower	Lygus bugs	1 pint per acre	Do not apply after flowering
Sorghum	Aphids	1 quart per acre	Do not apply within 21 days of harvest. Leaf injuries may occur on some varieties.
Spinach	Aphids	1 pint per acre	Do not apply within 14 days of harvest
		-or-	
		1 quart per acre	Do not apply within 21 days of harvest
Sugar beets	Armyworms	3/4 pint per acre	Do not apply within 20 days of harvest; 60 days if tops are to be fed to livestock
	Flea beetles		
	Leaf hoppers		
	Mites		
Sweet potatoes	Flea beetles	1½ pints per acre	Do not apply within 5 days of harvest
Tomatoes	Aphids	1 pint per acre	Do not apply within 10 days of harvest
	Armyworms		
	Flea beetles	-or-	
	Leafhoppers	1½ quarts per acre	Do not apply within 15 days of harvest
Turnips (foliage treatment)	Aphids	4/5 quart per acre	Do not apply within 15 days of harvest; 21 days if tops are to be used as food or feed.

For a mosquito larvicide on irrigated pastures, use 1/5 pint per acre up to 7 days of harvest.

NET CONTENTS

GALS.

1526-194
6-28-71
1842

ACCEPTED
JUN 28 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECOLOGICAL POISON REGISTER-
ED AND ~~NO~~ *286-194* SUBJECT
TO ATTACHED COMMENTS.

2042