



POISONOUS IF SWALLOWED

This product can kill you if swallowed, even in small amounts: spray mist or dust may be fatal if swallowed.

POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes.)


POISONOUS IF BREATHED

Breathing vapors, spray, mist or dust may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur.

Best Available COPY
POISON
 See Side Panels for Antidote & Precautions
 FLAMMABLE MIXTURE
 KEEP AWAY FROM HEAT OR OPEN FLAME



DANGER

Keep Out of Reach of Children

NOT FOR HOME USE

ARIZONA *Do not contaminate water by cleaning of equipment or disposal of wastes.*
 GROWERS' CHEMICAL CORPORATION
 Growing the Chemical Program
 Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 252-1111

METHYL PARATHION* EC-46
 EMULSIFIABLE
 Contains 4 pounds Methyl Parathion per gallon

ACTIVE INGREDIENTS:

O, O-dimethyl O-p-nitrophenyl thiophosphate	46%
Aromatic Petroleum solvent	46%
INERT INGREDIENTS:	6%
* Methyl Parathion	100%

For ground application dilute in 10 to 20 gallons of water per acre. For air application dilute in 3 to 10 gallons of water per acre.

CROP	INSECT PEST	MAXIMUM RATE	LIMITATION
ALFALFA (Calif-ornia)	aphids	3/4 pint per acre	Do not apply within 15 days of harvest
ALFALFA (States other than Calif.)	Armyworms Leafhoppers Mites	2 to 2-1/2 pints per acre	Do not apply within 15 days of harvest at 2 pint rate; 20 days above 2-1/2 pints rate
SMALL GRAINS (Barley, Oats, Rye, Wheat)	Aphids	1/2 pint per acre	No time limitations
	Armyworms Leaf hoppers	1-1/2 to 2-1/2 pints per acre	Do not apply within 15 days of harvest at 1-1/2 pint rate; Do not apply after heads begin to form at 2-1/2 pints

CROP	INSECT PEST	RATE	LIMITATIONS
Broccoli	Aphids	1 pint	Do not apply within 7 days
Brussels Sprouts	Aphids	1 pint	Do not apply within 20 days for each crop
Cabbage	Aphids	3 pints	Do not apply within 21 days of harvest
Carrots	Aphids Armyworms Fly beetles Leafhoppers	2 pints per acre	Do not apply within 15 days harvest. Do not use treated crop for food or feed
Cotton	Aphids Beehoppers Lycus bugs Thrips Cotton leaf perforator	3 quarts per acre	Do not apply within 5 days of harvest. Working in treated fields within 24 hrs. wear protective clothing.

This product is highly toxic and should be used only for direct treatment or residue control. Precautionary information may be obtained from your Cooperative Extension Service.

Best Available COPY

1526-194 2-7-72

10/24

Best Available COPY

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

WEAR SAFETY RULES

- (1) Wear water-proof pants, coat, hat, rubber boots or rubber overalls.
 - (2) Wear safety goggles.
 - (3) Wear mask or respirator approved by the U.S. Bureau of Mines for protection.
 - (4) Wear heavy duty natural rubber gloves.
- Keep all unprotected persons and children away from treated area or where there is danger of drift.
- Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to poison and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.
- IMPORTANT!** Before removing gloves, wash them with soap and water. Always wash hands, face and areas with soap and water before smoking, eating or drinking.
- AFTER WORK**, take off all work clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Resistor should be cleaned and filter replaced according to instructions included with respirator.

Red

POISON SIGNS (Symptoms)

Fazathion is a very dangerous poison. It readily enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment, or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, lightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

FIRST AID TREATMENT

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to poison and describe his condition.

If breathing has stopped, start artificial respiration immediately and maintain until doctor sees victim. If swallowed and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

NOTE TO PHYSICIAN

Antidote—administer atropine sulfate in large doses TWO TO FOUR mg. intravenously or intramuscularly as soon as diagnosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also indicated and may be given if atropine is not available.

ADOPHANT (M. IPAMQUIN) is a cholinesterase inhibitor and produces a strong cholinesterase inhibitor action on the central and peripheral nervous systems and producing pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and further relapses have been reported after initial improvement; VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding post-treatment areas.

CONTAINER DISPOSAL

Do not reuse empty drum. Return to drum conditioner or dealer by perforating or crushing and burying in a safe place.

NOTICE
Seller makes no warranty, expressed or implied, concerning the use of this product other than that indicated on the label. Buyer assumes all risk of use and/or handling of this material when used in a manner not intended or contrary to label instructions.

EPA Reg. No. 1526-19A
25547

NET CONTENTS 5 GALLONS

DATE

CROP	INSECT PER	MAXIMUM RATE	LIMITATIONS
Lettuces	Aphids Armyworms Leafhoppers Cabbage Looper	2 pints per acre	Do not apply within 21 days of harvest
Onions	Thrips	7/8 quart per acre	Do not apply within 15 days of harvest
Peas	Aphids Armyworms Flea beetles Leafhoppers	2 pints per acre	Do not apply within 5 days of harvest
Safflower	Lycus bugs	1 pint per acre	Do not apply after flowering
Sorghum	Aphids	2 pints per acre	Do not apply within 21 days of harvest. Seed treatment may reduce on some varieties.
Soybeans	Aphids	1 pint per acre	Do not apply within 21 days of harvest.

CROP	INSECT PER	RATE	LIMITATIONS
Sugar Beets	Armyworms Leaf beetles Cabbage Looper Moths	3/4 pint per acre	Do not apply within 20 days of harvest 60 days if from seed to be fed to livestock
Sweet Potatoes	White beetles Aphids Armyworms Cabbage Looper Moths	1 1/2 pints per acre	Do not apply within 5 days of harvest Do not apply within 10 days of harvest Do not apply within 15 days of harvest
Tomatoes	Aphids Armyworms Cabbage Looper Moths	1 pint per acre	Do not apply within 10 days of harvest Do not apply within 15 days of harvest
Turkey Corn (Cultivar)	Aphids	1 1/2 pints per acre	Do not apply within 15 days of harvest Do not apply within 21 days of harvest

Use only as directed on label. Do not use on crops not listed.

Best Available Copy

304

COLE CROPS (Broccoli, Brussels sprouts, Cabbage, Cauliflower)	Aphids	1 pint per acre	Do not apply within 10 days of harvest on Cabbages; 7 days on other Cole crops.
	Cabbage Looper	2 to 3 pints per acre	Do not apply within 21 days of harvest.

404

ACCEPTED
FEB 7 1972
UNDER THE FEDERAL INSECT
FUNGICIDE AND RODENTIDE ACT
FOR 2-15-72-1526-194
ED UNDER NO. 1526-194
TO ATTACHED COMMENTS. SUBJECT

1526-194