

UNICO SEVIN Zineb 3 - 5.2 Dust

CAUTION

Harmful if inhaled or swallowed. May cause irritation of nose and throat. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not contaminate feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid application during flowering or pollinating period when bees may be visiting the crop. If necessary to apply during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after the application.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke may be hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

To control the insects and the plant diseases on the field-grown crops listed below, apply at the rate specified for each crop, to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough, but do not apply excessive amounts. Begin applications when the insects or their damage first appear and before diseases appear (when disease is first reported in the area). Repeat as necessary, generally at 7-day intervals unless a shorter interval is stated. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES—Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall army worm, tarnished plant bug, Colorado potato beetle, early blight, late blight, leaf spot, and anthracnose on field grown tomatoes. Apply 40 to 50 pounds of dust per acre.

BEETS—Flea beetles, spittlebug, tarnished plant bug, stink bug, cabbage looper (when larvae are small; apply at 4 to 5-day intervals), and leaf spot. Apply 30 to 40 pounds of dust per acre. DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED: IF TOPS WILL NOT BE SO USED, DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

EGGPLANT AND PEPPERS—Leafhoppers, fall armyworm, European corn borer, tarnished plant bug, leaf spot, and phomopsis. Apply 30 to 40 pounds of dust per acre.

BEANS—Mexican bean beetle, bean leaf beetle, leafhoppers, rust, anthracnose and downy mildew on lima beans). Apply 30 to 40 pounds of dust per acre. DO NOT APPLY TO BEANS WITHIN 7 DAYS OF HARVEST.

CABBAGE, CAULIFLOWER, AND BROCCOLI—Harlequin bug, imported cabbage worm, larvae of diamond-back moth, cabbage looper (when larvae are small; apply at 4 to 5-day intervals), and downy mildew. Apply 30 to 40 pounds of dust per acre. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

CUCUMBERS, PUMPKINS, AND SQUASH—Pickle worm, melon worm, cucumber beetles, flea beetles, leafhoppers, squash bug, downy mildew, Ascochyta blight (on cucumbers), and anthracnose (on squash). Apply 30 to 33 pounds of dust per acre.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

ACCEPTED