

**ANTIDOTE AND FIRST AID STATEMENT**

**CAUTION**

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Do not get in eyes. Avoid prolonged or repeated breathing of dust. Skin contact may be harmful. Avoid contact. If spilled on skin, wash thoroughly with soap and water. Wash hands before eating. Take shower or bath after work. Do not contaminate feed or foodstuffs. Wear regular long-sleeve work clothing. Change to clean clothing daily.

**ANTIDOTE AND FIRST AID**

Give an emetic of warm mustard water - CALL A PHYSICIAN. Take one or two 1/100 grain atropine tablets and repeat in 1 hour if necessary.

**CAUTION**

Do not graze treated fields or feed trash to dairy or meat animals. Use only for recommended purpose and at recommended dosages.

Do not reuse empty container. Destroy it by burying or burning. Stay out of smoke and fumes.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**BEE CAUTION**


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service

**STOP - READ LABEL BEFORE USING!**

**Agro-Chem Brand**

**CHEMICAL DISTRIBUTORS**

**COBA**



**P.O. BOX 21537 PHOENIX, ARIZ. 85016 PHONE 602 244-1311**

**E.P.A. REG. NO. 1516-1-64  
E.P.A. EST. NO. 1526-AZ-1**

**SEVIN 7.5 with 50% SULFUR**

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl-N-methylcarbamate)----- 7.50  
Sulfur----- 50.00%

**INERT INGREDIENTS:----- 42.50%**  
**100.00%**

\*Trade-mark of Union Carbide Corporation

DO NOT APPLY THIS MATERIAL WITH EQUIPMENT THAT HAS EVER BEEN USED TO APPLY 2,4-D, 1,4,5-T OR RELATED WEED KILLING COMPOUNDS.

Do not allow to drift to sulfur allergic crops such as cucurbits (melons, squash, cucumbers, etc.) and lettuce.

This product is for use on hopper, cotton, pink bollworm, etc. For control of cotton. For control of insects, apply at the rate of 10 to 26 pounds per acre. For control of insects, begin application as soon as necessary. For control of insects, apply further as necessary.

Net CONTENTS



**CAUTION: Keep Out of Reach of Children.**

See other precautions on this label

