

**DANGER: Keep Out of Reach of Children: Harmful if Swallowed. Do Not Inhale**

WARNING

May be fatal if swallowed, inhaled or absorbed thru skin! Rapidly absorbed thru skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Methyl Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

ANTIDOTE


IF SWALLOWED: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

CAUTION

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be harmed.

Do not reuse container. Destroy when empty.




U. S. D. A. Reg. No. 1526-337

**ACTIVE INGREDIENTS:**

0,0-dimethyl, 0-p-nitrophenyl thiophosphate*	2.00%
Sulfur	80.00%

<b>INERT INGREDIENTS:</b>	<b>18.00%</b>
	<b>100.00%</b>

\*Methyl Parathion



Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

DIRECTIONS FOR USE

This insecticide is for the control of aphids, red spiders, beet armyworm, yellow-striped armyworm, cabbage loopers, alfalfa semi-looper, lygus, thrips, and grasshoppers on alfalfa, carrots, barley and wheat. Apply at a rate of 15 to 35 pounds per acre depending upon the type of equipment used, the size and growth of the crops, and the insect to be controlled. Further dustings should be governed by populations of damaging insects. For further information consult your state agricultural experiment station. Do not apply within 15 days of harvest to any of the crops.

CAUTION

Do not apply when weather conditions favor drift from areas treated. Do not allow to drift to sulfur allergic crops such as cucurbits (melons, squash, cucumbers, etc.) or to lettuce.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**DANGER POISON**

**DANGER KEEP OUT OF REACH OF CHILDREN SEE WARNING AND ANTIDOTE STATEMENTS ON SIDE PANEL**

NET CONTENTS

LBS.