

DANGER: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale.

will change on next printing
WARNING

May be fatal if swallowed, inhaled or absorbed thru skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not re-use container. Destroy by burying in a safe place. Do not burn.

ANTIDOTE


If swallowed: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY.

CAUTION

Do not enter treated fields within 24 hours after application. Should wear protective clothing. Use only for intended purposes at recommended dosages. Apply this product only as specified on this label. Do not apply when weather conditions favor drift to areas treated.

CAUTION

Do not feed birds, sheep or calves, etc. *Birds feeding on treated areas may be killed.*



U. S. D. A. Reg. No. 1526-953

ACTIVE INGREDIENTS:

Parathion*	2.00%
Endosulfan (Hexachlorohexahydro-methano-2,4,3-benzodioxathiepin oxide)	3.00%
INERT INGREDIENTS:	95.00%
	100.00%

(R) Trademark of Farbwerke Hoechst A.C.
 U.S. Patent No. 2,799,685

* O,O-diethyl O-p-nitrophenyl thiophosphate

CAUTION

DO NOT APPLY THIS MATERIAL WITH EQUIPMENT THAT HAS EVER BEEN USED TO APPLY 2,4-D, 2,4,5-T OR RELATED KILLING COMPOUNDS.

*9-25-70
 1526-353*



DANGER POISON



Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

DIRECTIONS FOR USE

CROP	INSECT PESTS	MAXIMUM RATES	LIMITATIONS
Broccoli	Cabbage Aphids	25 lbs. per acre	Do not apply within 10 days of harvest. Do not apply after edible parts start to form. Do not apply within 14 days of harvest on cauliflower.
Cabbage	Cabbage Looper		
Cauliflower	Imported Cabbage Worm		
	Diamond Back Moth Larvae		
	Flea Beetle		
Cotton	Boll Weevil	50 lbs. per acre	Do not apply after bolls open. Do not graze meat or dairy animals in treated fields. Do not apply within 5 days of hand picking.
	Cotton Stink Bug		
	Bugs		
	Lygus Bug		
	Boll Worm		
	Cabbage Looper		
	Cotton Leaf Perforator		
Grapes	Grape Leafhopper	50 lbs. per acre	Do not apply within 14 days of harvest.
Lettuce	Cabbage Looper	25 lbs. per acre	Do not apply more than 3 times after thinning. Do not apply within 21 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.
	Green Peach Aphid		
	Diamond Back Moth larvae		
	Imported Cabbage Worm		

(Continued)

NOTICE

Seller makes no warranty, express, implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling is contrary to label instructions.

CROP	INSECT PESTS	MAXIMUM RATES	LIMITATIONS
Melons	Aphids	25 lbs. per acre	Do not apply within 7 days of harvest.
	Cucumber Beetles		
	Cabbage Looper		
	Omnivorous Leaf Roller		
Potatoes	Flea Beetle	<i>50</i> lbs. per acre	Do not apply within 5 days of harvest.
	Colorado Potato Beetle		
	Leafhopper		
	Aphids		
	Green Stink Bugs		
Tomatoes	Potato Psyllid		
	Hornworm	33 lbs. per acre	Do not apply within 10 days of harvest.
	Tomato Fruitworm		

REGISTERED TRADEMARK OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

NET WEIGHT

LBS.