

WARNING & ANTIDOTE STATEMENT

WARNING

May be Fatal if Swallowed, Inhaled, or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Keep all unprotected persons out of operating areas where there may be danger of drift.

ANTIDOTE

If Swallowed: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a Physician immediately!

If on Skin: In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

PHOENIX BRAND PARATHION 2 WITH 80% SULFUR

U. S. D. A. Reg. No. 1526—368

ACTIVE INGREDIENTS:

Parathion-----	2.00%
Sulfur-----	80.00%
INERT INGREDIENTS:-----	18.00%
	100.00%

*0,0-diethyl,0-p-nitrophenyl thiophosphat₄


ACCEPTED

OCT 16 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
AND ECONOMIC POISON REGISTERED
UNDER NO. 126365 SUBJECT
TO ATTACHED COMMENTS.



DANGER POISON



Main Office 734 East S. P. Drive -- Phoenix, Arizona 85001 -- Phone 253 1111

DIRECTIONS FOR USE

This insecticide is for the control of aphids, red spiders, beet armyworm, yellow-striped armyworm, loopers, lygus, thrips and grasshoppers on alfalfa, carrots, barley and wheat. Apply at a rate of 15 to 20 pounds per acre depending upon the type of equipment used, the size and growth of the crops, and the insect to be controlled. Further dustings should be governed by populations of damaging insects. For further information on this material, consult your state agricultural experiment station. Do not apply within 15 days of harvest to any of the crops.

WARNING

Do not allow to drift to sulfur allergic crops such as cucurbits (melons, squash, cucumbers, etc.) or to lettuce.

Do not reuse containers. Destroy by burying in a safe place. Do not burn. Do not contaminate water by cleaning of equipment.

This product toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

NOTICE

Seller makes no warranty, expressed or implied, as to the safety of use of this product other than indicated on the label. There is a known risk of use and/or handling of this material which use and/or handling is contrary to label instructions.