

CAUTION: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale

CAUTION

Harmful if taken internally. Avoid inhalation of dust. Avoid contact with skin, eyes, or clothing. Keep away from fire and sparks.

Keep product dry to prevent reduced effectiveness and fire hazard. Use this product only for recommended purposes at recommended dosages.


ANTIDOTE

If on skin flush with plenty of water. If in eyes, flush with water and call a physician.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ACCEPTED
3/15/72
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1526-373



EPA Reg. No. 1526-373

ACTIVE INGREDIENTS:

Zineb (Zinc Ethy ene Bis
Dithiocarbamate)----- 5.20%

INERT INGREDIENTS:-----94.80%
100.00%

ZINEB 5 FUNGICIDE dust may be applied to moist or dry foliage anytime dusting conditions are favorable.

Begin application at first sign of or prior to appearance of disease and repeat as indicated.

DIRECTIONS FOR USE

CROP	PLANT DISEASES	MAXIMUM RATE	LIMITATIONS
Tomatoes	Early & late blight Anthracnose Stemphylium Septoria Leaf Spots	73 lbs. per acre	Apply at 7 to 10 day intervals. No time limitations.
Wheat	Stem & Leaf Rust	43 lbs. per acre	Do not apply within 3 to 4 weeks (to milk stage) of harvest.

ACCEPTED
3/15/72
1526-373

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED

CAUTION: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale. SEE WARNING AND ANTIDOTE STATEMENTS ON SIDE PANEL

ACCEPTED
3/15/72

1526-373
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

ARIZONA
GRO-CHEMICAL
CORPORATION

Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

DIRECTIONS FOR USE

CROP	PLANT DISEASES	MAXIMUM RATE	LIMITATIONS
Broccoli	Downy Mildew	92 lbs. per acre	Apply at 7 to 14 day intervals to within 7 days of harvest
Cabbage	Alternaria		
Carrot	Leaf Blight Cercospora Leafspot	57 lbs. per acre	Apply at 7 to 14 day intervals to within 7 days of harvest if treated tops are to be used as food or feed.
Cantaloupe	Downy mildew Alternaria Anthracnose	73 lbs. per acre	Apply at 7 to 14 day intervals. No time limitations.
Grapes	Downy mildew Black rot Ripe rot	57 lbs. per acre	Apply just before and just after bloom and again 10 days. Do not apply within 7 days of harvest.
Lettuce	Downy mildew	57 lbs. per acre	Apply at weekly intervals to within 5 days of harvest on head lettuce and 7 days on Leaf Lettuce.
Onions	Botrytis Leaf Blight Purple Blotch	57 lbs. per acre	Apply where plants reach 4 leaf stage, and continue at 7 to 10 day intervals up to within 7 days of harvest on green onions.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and of handling of this material when such use and handling is contrary to label instructions.

NET CONTENTS

CAUTION
 Harmful if taken internally. Avoid inhalation or
 dust. Avoid contact with skin, eyes, or clothing.
 Keep away from fire and sparks.

Keep product dry to prevent reduced effectiveness
 and fire hazard. Use this product only for re-
 commended purposes at recommended dosages.

ANTIDOTE
 If on skin flush with plenty of water. If in
 eyes, flush with water and call a physician.

CONTAINER DISPOSAL
 Do not reuse empty container. Destroy it by
 burying with waste or burning. Stay away from
 smoke or fumes.

ACCEPTED
 3/15/72
 1526-373

ZINEB 5 FUNGICIDE

ZINEB 5
 FUNGICIDE
 EPA Reg. No. 1526-373

ACTIVE INGREDIENTS:
 Zineb (Zinc Ethylene Bis-
 Dithiocarbamate)----- 5.20%
INERT INGREDIENTS:----- 94.80%
 100.00%

ZINEB 5 FUNGICIDE may be applied to moist
 or dry foliage anytime dusting conditions are
 favorable.

Begin application at first sign of or prior to
 appearance of disease and repeat as indicated.

DIRECTIONS FOR USE

CROP	PLANT DISEASES	MAXIMUM RATE	LIMITATIONS
Tomatoes	Early & late blight Anthracnose Stemphylium Septoria Leaf Spots	73 lbs. per acre	Apply at 7 to 10 day intervals. No time limitations.
Wheat	Stem & Leaf Rust	43 lbs. per acre	Do not apply within 3 to 4 weeks (from milk stage) of harvest.

ACCEPTED
 3/15/72
 1526-373

KEEP OUT OF REACH OF CHILDREN
 READ THE DIRECTIONS, WARNINGS AND ANTIDOTE STATEMENTS ON THE LABEL

ACCEPTED
 3/15/72
 1526-373

CROP	PLANT DISEASES	MAXIMUM RATE	LIMITATIONS
Broccoli	Downy Mildew	92 lbs. per acre	Apply at 7 to 14 day intervals to within 7 days of harvest
Cabbage	Alternaria		
Carrot	Leaf Blight Cercospora Leafspot	57 lbs. per acre	Apply at 7 to 14 day intervals to within 7 days of harvest if treated tops are to be used as food or feed
Cantaloupe	Downy mildew Alternaria Anthracnose	73 lbs. per acre	Apply at 7 to 14 day intervals. No time limitations.
Grapes	Downy mildew Black rot Ripe rot	57 lbs. per acre	Apply just before and just after bloom and again 10 days. Do not apply within 7 days of harvest.
Lettuce	Downy mildew	57 lbs. per acre	Apply at weekly intervals to within 5 days of harvest on head lettuce and 7 days on leaf lettuce.
Onions	Botrytis leaf blight Purple Blotch	57 lbs. per acre	Apply before plants reach 4 leaf stage, and continue at 7 to 10 day intervals up to within 7 days of harvest on green onions.