

CAUTION

Keep Out of Reach of Children



**CAPTAN 10 with
50% SULPHUR**

ACTIVE INGREDIENTS:	
Captan	10.00%
Sulphur	50.00%
INERT INGREDIENTS: 40.00%	
USDA REG. NO. 1526 100.00%	

WARNING

Harmful if swallowed, inhaled or absorbed thru skin. Do not take orally. If spilled on skin or clothing, wash thoroughly with soap and water. Wash clothing with soap and water before re-use. Do not contaminate feed and foodstuffs. Keep out of reach of children and domestic animals. Wash hands before eating or smoking.

ANTIDOTE AND FIRST AID

Give an emetic of warm mustard water. CALL A PHYSICIAN.

DIRECTIONS FOR USE

This product is for the control of Bunch rot on grapes. Apply at the rate of 35 to 50 lbs. per acre in a manner to give thorough coverage of the affected portions of the crop plants. Begin applications when a damageable or potentially damageable infestation of the rot appears, and continue treatments at weekly intervals as additional control is needed. For further specific information, consult your local agricultural authorities.

Handwritten signatures and initials on the left side of the label.

ACCEPTED
Apr 21, 1967
1526-399
Comments

CAUTION

Keep Out of Reach of Children

POISON

PHOENIX BRAND

**CAPTAN 10 with
50% SULPHUR**

ACTIVE INGREDIENTS:

Captan *10* 50.00%
Sulphur *50* 50.00%

INERT INGREDIENTS: 40.00%

USDA REG. NO. 1526 100.00%

WARNING

Harmful if swallowed, inhaled or absorbed thru skin. Do not take internally. If spilled on skin or clothing, wash thoroughly with soap and water. Wash clothing with soap and water before re-use. Do not contaminate feed and foodstuffs. Keep out of reach of children and domestic animals. Wash hands before eating or smoking.

ANTIDOTE AND FIRST AID

Give an emetic of warm mustard water. CALL A PHYSICIAN.

DIRECTIONS FOR USE

This product is for the control of Bunch rot on grapes. Apply at the rate of 35 to 50 lbs. per acre in a manner to give thorough coverage of the affected portions of the crop plants. Begin applications when a damageable or potentially damageable infestation of the rot appears, and continue treatments at weekly intervals as additional control is needed. For further specific information, consult your local agricultural authorities.

Handwritten signature and scribbles on the left side of the label.

ACCEPTED
Apr. 21, 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
REGISTERED
UNDER NO. 1526-399
Comments

3M 5-63

HIGHLY TOXIC TO FISH AND WILDLIFE. DO NOT CONTAMINATE ANY BODY OF WATER OR APPLY TO ANY AREA NOT SPECIFIED ON THIS LABEL.

CAUTION

~~To protect fish and wildlife, do not contaminate streams, lakes or ponds with this dust. Use only for recommended purposes and at recommended dosages.~~
Do not allow this dust to drift onto sulphur allergic cucurbits (melons, cucumbers, squash, etc.) and lettuce crops.

DO NOT APPLY THIS MATERIAL WITH EQUIPMENT THAT HAS EVER BEEN USED TO APPLY 2,4-D, 2,4,5-T OR RELATED WEED-KILLING COMPOUNDS.

NET WEIGHT 50 LBS.

Manufactured in Arizona by:



734 East S. P. Drive - Phone 253-1111
P. O. Box 2191, Phoenix, Arizona

Manufacturing and Sales facilities in
PHOENIX - CHANDLER - TOLTEC - YUMA - WILLCOX

ACCEPTED WITH COMMENTS

3M 5-63

HIGHLY TOXIC TO FISH AND WILDLIFE. DO NOT CONTAMINATE ANY BODY OF WATER NOR APPLY TO ANY AREA NOT SPECIFIED ON THIS LABEL.

CAUTION

~~To protect fish and wildlife, do not contaminate streams, lakes or ponds with this dust. Use only for recommended purposes and at recommended dosages.~~
Do not allow this dust to drift onto sulphur allergic cucurbits (melons, cucumbers, squash, etc.) and lettuce crops

DO NOT APPLY THIS MATERIAL WITH EQUIPMENT THAT HAS EVER BEEN USED TO APPLY 2,4-D, 2,4,5-T OR RELATED WEED-KILLING COMPOUNDS.

NET WEIGHT 50 LBS.

Manufactured in Arizona by:



734 East S. P. Drive - Phone 253-1111
P. O. Box 2191, Phoenix, Arizona

Manufacturing and Sales facilities in
PHOENIX - CHANDLER - TOLTEC - YUMA - WILLCOX

ACCEPTED WITH COMMENTS

Handwritten scribbles on the left margin.