STOP READ LABEL BEFORE USING

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

DBA



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526-405 EST. NO. 1526-AZ-1

D-LEAF-M

cotton defoliant

ghly with water

kin or clothing.

ghly after using

feed, or other

d areas. Do not

n to livestock.

reams. se only recom-

D-Leaf M'is a liquid defoliant containing 2 pounds of Sodium Chlorate per gallon at 60°F.

**ACTIVE INGREDIENT(S):** 

 DIRECTIONS FOR USE

For best results D-Leaf M should be applied to mature plants. The usual signs of maturity include a slight yellowing of the foliage and a reddish tinge on the leaves and stems. If there is any question as to the maturity of the plants, consult your state agricultural experiment station.

Plants should not be stressed for water when D-Leaf M is applied since tough wilted leaves tend to hinder the uptake of the chemical and prevent growth of the abscission layer.

D-Leaf M should be applied at the rate of 2 gallons per acre. If an airplane applicator is used the 2 gallons should be diluted with 8 gallons of water.

If a ground applicator is used the 2 gallons should be diluted with from 15 to 30 gallons of water. Do not apply within 7 days of harvest.

D-Leaf M is not selective in action. Care should be exercised in applying near desirable plants. Avoid drift or contact with plants other than those being treated. Do not apply in windy weather.

NOTICE

## STATEMEN

## Draiin agro-unem CHEMICAL DISTRIBUTORS DBA

RIZONA GROCHEMICAL COMPANY owing thru Chemical Progress

P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526-405 EST. NO. 1526-AZ-1

D-LEAF-M

cotton defoliant

oughly with water

i skin or clothing

oughly after using

d. feed, or other

ated areas. Do not

ton to livestock.

Use only

streams.

recom-

D-Leaf M is a liquid defoliant containing 2 pounds of Sodium Chlorate per gallon at 60°F.

ACTIVE INGREDIENT(S):

100.00%

For best results D-Leaf M should be applied to mature plants. The usual signs of maturity include a slight yellowing of the foliage and a reddish tinge on the leaves and stems. If there is any question as to the maturity of the plants, consult your state agricultural experiment station.

Plants should not be stressed for water when D-Leaf M is applied since tough wilted leaves tend to hinder the uptake of the chemical and prevent growth of the abscission layer.

D-Leaf M should be applied at the rate of 2 gallons per acre. If an airplane applicator is used the 2 gallons should be diluted with 8 gallons of water.

If a ground applicator is used the 2 gallons should be diluted with from 15 to 30 gallons of water. Do not apply within 7 days of harvest

D-Leaf M is not selective in action. Care should be exercised in applying near desirable plants. Avoid drift or contact with plants other than those being treated Do not apply in windy weather.

CAUTION: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale.

