Main Office: 734 East S. P. Drive --- Phoenix, Arizona 85001 --- Phone 253-1111

DIRECTIONS FOR USE

This insecticide is for the control of aphids, cabbage looper and thrips on spinach, imported cabbage worm and cabbage looper on broccoli, cabbage and caulitlower. On potatoes, this insecticide is for the control of aphids, flea beetle, leafhopper and white flies. This material is also a fungicide used to prevent and control downy mildew of spinach and Brussels sprouts. Begin application at the first indication of any of the diseases (especially when the humidity is favorable for the spread of the disease) at a rate of 25 pounds per acre in a manner to insure thorough coverage of the affected portions of the crop-plants. This product must not be used unless residues are removed by stripping, trimming and washing. For further specific information on the use of this product, especially as to unusual weather, climate, or soil conditions, consult your state agricultural extension service. Apply in a manner to give thorough coverage of the affected portions of the crop-plants when a damageable or potentially damageable infestation of the insectpests occur. Continue treatments as additional control is needed, heeding the following dosages and limitations.

BROCCOLI: Use 13 pounds per acre. Do not apply within 7 days of harvest.

BRUSSEL SPROUTS: Use 25 pounds per acre. Do not apply within 7 days of harvest.

CABBAGE: Use 25 lbs. per acre. Do not apply within 7 days of harvest.

CAULIFLOWER: Use 25 pounds per acre. Do not apply after edible parts start to form.

SPINACH: Apply at 25 pounds per acre. Do not apply within 14 days of harvest.

POTATOES: Use 30-35 pounds per acre. Apply when insects first appear and repeat as required. Do not apply within 5 days of harvest.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

ाम (क्रामाराक

remaining addition of the Mayder of

WARNING: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale.

WARNING

May be fatal if swallowed, inhaled or absorbed thru skin! RAPIDLY ABSORBED THRU SKIN! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

WARNING

Workers entering treated fields within 24 hours after application should wear protective clothing. Keep away from fire and sparks.

ANTIDOTES

If swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately!

If on skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

PEQUENIX BRAND

PARATHION 2 MANEB 5

U. S. D. A. Reg. No. 1526-410

ACTIVE INGREDIENTS:	2 00%
Parathion*	2.00%
Maneb (Manganous Ethylene	
Bisdithiocarbamate)	5.00%
INERT INGREDIENTS:	
	100.00%
*0,0-Diethyl O-p-nitrophenyl Thiophosphate	

CAULION

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on label. Destroy container by burying in a safe place. Never reuse. Use only for recommended purposes at recommended dosages.

Do not apply this material with equipment that has ever been used to apply 2,4-D, 2,4,5-T or related weed killing compounds.

