

WARNING: Keep Out of Reach of Children - Harmful if Swallowed - Do Not Inhale.

WARNING AND ANTIDOTE STATEMENTS

WARNING

Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. When container is empty, wash thoroughly and destroy. Never re-use. Stay away from smoke and fumes.

ANTIDOTES

IF SWALLOWED. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a Physician Immediately!


CAUTION

Keep out of reach of children and domestic animals. Do not contaminate feed or foodstuffs. Use only for recommended purposes at recommended dosages.

CAUTION

Toxic to fish. Keep out of lakes, streams, or ponds.

12-20-69
1526-428



U. S. D. A. Reg. No. 1526-428

ACTIVE INGREDIENTS:
Malathion*----- 5.00%
INERT INGREDIENTS:----- 95.00%
100.00%

*O,O-dimethyl Dithiophosphate of diethyl mercaptosuccinate.

WARNING

KEEP OUT OF REACH OF CHILDREN

LARGER TYPE

WARNING: Keep Out of Reach of Children - Harmful if Swallowed - Do Not Inhale. SEE WARNING AND ANTIDOTE STATEMENTS ON SIDE PAN

RIGHT PANEL



Main Office: 734 East S. P. Drive - Phoenix, Arizona 85001 - Phone 253-1111

Crop	Insect Pest	Maximum Rate	Limitations
Alfalfa	Pea aphid Spotted alfalfa aphid Cowpea aphid	25 to 30 lbs. per acre	Do not apply within 7 days of harvest. Do not apply during full bloom.
Broccoli Brussels sprouts Cabbage	Cabbage aphid Turnip aphid	20 to 50 lbs. per acre	Do not apply within 7 days of harvest.
Cotton	Cotton aphid White flies Southern garden leafhopper Lygus Cotton leaf perforator	25 to 50 lbs. per acre	No time limitations.
Grapes	Leafhoppers Aphids	20 to 50 lbs. per acre	Do not apply within 3 days of harvest.
Lettuce (head)	Green peach aphid Barr's aphid Thrips	20 to 50 lbs. per acre	Do not apply within 7 days of harvest.
Potatoes	Green peach aphid Potato aphid Leafhoppers	25 to 50 lbs. per acre	No time limitation

For further specific information, consult your State Agricultural Extension Service.

NOTICE
Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NET CONTENTS 1.5 GALS.

STOP-READ LABEL BEFORE USING!

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

D.B.A.



Agro-Chem Brand Progress

P.O. BOX 17, HOUSTON, TEXAS 77001

F.P. REG. NO. 1125-428
EST. NO. 1925-AZ-1

Agro-Chem

WARNING

Avoid contact with skin. Avoid prolonged breathing of dust or mist. Avoid prolonged or repeated contact with skin. Wash thoroughly after use. Avoid contamination of food and feedstuffs. Keep out of reach of children. When contact with skin occurs, wash thoroughly and frequently. Do not use in a safe place. Do not use. Stay away from smoke and flames.

IF SWALLOWED: Do not induce vomiting. Give water to drink and keep patient cool until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately!

Keep out of reach of children and domestic animals. Do not contaminate feed or feedstuffs. Use only for recommended purposes at recommended dosages.

Toxic to fish. Keep out of lakes, streams, or ponds.

This dust should be applied in a manner to give thorough coverage of the affected portions of the crop-plants. Begin applications when a damageable or potentially damageable infestation of the insect pests appear and continue treatments as control is needed.

MALATHION 5 DUST

MALATHION

ACTIVE INGREDIENTS:

Malathion*	5.00%
INERT INGREDIENTS	95.00%
	<u>100.00%</u>

* O, O-dimethyl Dithiophosphate of diethyl mercaptosuccinate

Do not use this product for any uses not

NOTE: Important - Do not allow dry application would be closer to harvest be applied as a foliar application only.

Camerman's Note
Poor Copy

STOP-READ LABEL BEFORE USING

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

DRA



AGRO-CHEM BRAND
DISTRIBUTORS
P.O. BOX 17
MORONG, PANGASINAN
PHILIPPINES

P.O. BOX 17 MORONG, PANGASINAN PHILIPPINES 11311

F.P. REG. NO. 1125-128
EST. NO. 1125-AZ-1

MALATHION 5 DUST

ACTIVE INGREDIENTS:

Malathion*	5.00%
INERT INGREDIENTS	95.00%
		<u>100.00%</u>

* O, O-dimethyl Dithiophosphate of diethyl mercaptosuccinate

Camerman's Note
Poor Copy

Agro-Chem Brand

MALATHION 5 DUST

Do not use this product for any uses other than those specified on this label.

NOTE: Important -- Do not allow drift of this product to crops for which its use is not registered or to crops for which its use may be registered but application would be closer to harvest than recommended. This product is to be applied as a foliar application only.

TEAR ALONG THIS LINE

PLANT	PLANT PEST	DOSAGE RATE: USE	LIMITATIONS
Brussels Sprouts, Cauliflower	Caterpillars	5 lbs.	Do not apply within 3 days before harvest.
Bentlounge	Cucumber beetles	25 lbs.	Do not apply within 1 day before harvest.
Clover	Aphids, mites	25 - 30 lbs.	None
Collards	Caterpillars	25 lbs.	Do not apply within 7 days before harvest.
Cotton	Fall Armyworms, garden webworms	25 - 50 lbs.	None
	Fleahoppers	10 - 20 lbs.	
Cucumbers	Cucumber beetles	35 lbs.	Do not apply within 1 day before harvest.
Grass, grass hay	Aphids, leafhoppers, Armyworms	30 - 40 lbs. 30 - 35 lbs.	None
Lettuce	Leafhoppers	25 lbs.	Leaf lettuce — Do not apply within 14 days before harvest. Head Lettuce — Do not apply within 7 days before harvest.
Mint	Adult flea beetles, caterpillars	20 lbs.	Do not apply within 7 days before harvest.
Mustard greens	Flea beetles	20 - 25 lbs.	Do not apply within 7 days before harvest.
Onion	Aphids	15 - 25 lbs.	Do not apply within 1 day before harvest.
	Japanese beetles	25 lbs.	
Onions	Onion maggots	30 lbs.	Do not apply within 3 days before harvest.
Pasture, range grass	Aphids, leafhoppers	30 - 40 lbs.	None.
Peanuts	Leafhoppers	20 lbs.	None.
Peas	Aphids, pea weevils	20 - 25 lbs.	Do not apply within 3 days before harvest. If vines are to be fed, do not apply within 7 days before harvest.
Pecans	Pecan nut casebearers	80 lbs.	None
Peppers	Maggots	30 lbs.	Do not apply within 3 days before harvest.
Potatoes	False chinch bugs	20 lbs.	None
	Mealybugs	30 lbs.	
Pumpkins	Squash vine borers	35 lbs.	Do not apply within 3 days before harvest. Do not apply when plants are wet.
Soybeans	Mexican bean beetles	30 - 40 lbs.	Do not apply within 3 days before harvest.
Squash	Cucumber beetles, squash vine borers	35 lbs.	Do not apply within 1 day before harvest.
Strawberries	Field crickets, thrips	20 - 40 lbs.	Do not apply within 3 days before harvest.

PEST	DOSAGE RATE: USE	LIMITATIONS
Millipedes, sowbugs, springtails	1 lb./150 square feet of soil surface or where insects congregate. Repeat at 7 - 10 day intervals or as needed.	None.

FOR USE ON LIVESTOCK:			
ANIMAL	ANIMAL PEST	DOSAGE RATE: USE	LIMITATIONS
Beef and non-milking cattle, horses	Horn flies	3 tablespoons per animal. Apply on back and neck of animal and repeat at 10 - 14 day intervals.	Do not apply to non-lactating dairy animals within 2 weeks of freshening. Do not apply to animals under 1 month of age. May be applied on day of slaughter.
	Ticks	Apply to back and neck of animals.	
	Horn flies, ticks, and in reduction of face flies.	For use in dust bag self applicators only. 10-20 lbs./applicator. Arrange dust bags at mid shoulder height. Do not contaminate salt, feed, or water.	
Dairy cattle	Horn flies	3 tablespoons/animal. Apply on back and neck of animals and repeat at 10 - 14 day intervals.	Do not apply to dairy cattle within 5 hours before milking or during milking.
Sheep, Goats	Keds, lice, ticks	Treat animals thoroughly. Apply with hand or power duster. Repeat in 2 - 3 weeks if needed.	Do not apply to milk goats. Do not apply to animals under 1 month of age.
Hogs	Lice	Adult hogs - apply thoroughly to animals and pens. Animals under 1 month of age - Apply 1/4 - 1/2 tablespoon per animal. Repeat only as necessary.	Avoid contamination of feed, food containers, and watering troughs.
Chickens, Ducks, Geese, Turkeys	Chicken red mites, lice, northern fowl mites.	Dust individual birds, use shaker can or rotary hand duster only. 1 lb. / 50-60 square feet premise treatment only. Apply to litter, walls, ceilings, roosts, nests, and adjacent areas. Force into cracks and crevices. Use hand duster or puff duster.	Do not contaminate feed, feeders, or water.
	Chiggers	20 lbs./acre. Range treatment only. Apply 1 day before placing poultry on range. Repeat every 2 - 3 weeks.	
	Lice, northern fowl mites	For use in dust bath boxes only. 1 lb. box (18"x12"x3"/30 hens).	

Out of Reach of Children
 Read instructions on this label.

ORSA

TOP-LEAD LABEL FOR USING:

ANIMAL	ANIMAL PEST	DOSAGE RATE: USE	LIMITATIONS
Chickens, Ducks, Geese, Turkeys, Con't.	Stick tight fleas	For use in dust bath boxes only. 12 lbs. box (2'x3'x1'/100 birds). For use in natural soil walls only. 6 lbs. / 50 birds. For use in range pens only. 3 1/2 lbs. / 20 square feet. Repeat in 28 days and then as necessary. For use in brooder houses only. 1 1/2 lbs. 100 chicks. Distribute dust evenly in each corner of brooder house.	Do not contaminate feed, feeders, or water.

FOR USE ON OUTDOOR ORNAMENTALS:			
PLANT	PLANT PEST	DOSAGE RATE: USE	LIMITATIONS
Flowers, shade trees, shrubs.	Aphids, spider mites, bagworms.	Apply for thorough coverage.	None.