

## WARNING & ANTIDOTE STATEMENT

### ANTIDOTE

If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY! If on Skin, remove contaminated clothing and immediately wash skin with soap and water. NOTE TO PHYSICIANS: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

### WARNING

Poisonous if swallowed, inhaled or absorbed through skin! Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Phosphamidon protection. Rapidly absorbed through skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, water proof protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and hot water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. DO NOT APPLY UNDER CONDITIONS INVOLVING POSSIBLE DRIFT TO FOOD, FORAGE OR OTHER PLANTINGS THAT MIGHT BE DAMAGED OR THE CROPS THEREOF RENDERED UNFIT FOR SALE, USE OR CONSUMPTION.

WHEN CONTAINER IS EMPTY, IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER RE-USE.

### CAUTION

This product is toxic to fish, shrimp, crabs, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Avoid use during flowering and pollination periods. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**STORE IN COOL DRY PLACE  
DO NOT TAKE INTERNALLY  
DO NOT INHALE — DO NOT GET ON SKIN**

STOP-READ LABEL BEFORE USING!

## Agro-Chem Brand

CHEMICAL DISTRIBUTORS

D B A



P. O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526-431A

E.P.A. EST. NO. 1526-AZ-1

# PHOSPHAMIDON 8 SPRAY


### ACTIVE INGREDIENTS:

Phosphamidon (2-chloro-2-diethylcarbamoyl-1-methylvinyl dimethyl phosphate)	74.5%
Related Compounds	3.5%
INERT INGREDIENTS:	22.0%
	100.0%

Contains 8 lbs. Phosphamidon Insecticide per Gallon.  
U.S. Patent No. 2,908,605

PHOSPHAMIDON 8 SPRAY is a liquid systemic insecticide for the control of both sucking and chewing insects. It is intended to be applied as a diluted spray and is suitable for application by ground or air equipment.

**POISON**



**DANGER**

## DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

Keep pesticide in original container. Do not use food or drink containers for diluting.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:  
GROUND APPLICATION — Vegetable and Field Crops — 30 to 250 gals. Deciduous Fruit Crops — 60 to 800 gals. Citrus — 400 to 3,000 gals. (On young Fruit Plantings — Use a minimum of 15 gals. water per acre).

Where Recommended for AIR APPLICATION — Field Crops — 3 to 10 gals. Orchard Crops — 10 to 20 gals.

NOTE: Important — Avoid drift of this product to crops for which it is not registered or to any crop for which its use may be registered but application would be closer to harvest than recommended. This product is to be applied as a foliar application only.

CROP	CROP PEST	DOSAGE USE	RATE	LIMITATIONS
Apples	Aphids	1/6 gal	pint/100 water	Do not apply within 30 days before harvest. Do not apply more than 1 quart of this product per acre.

Green Peach Aphids	1/4 gal	pint/100 water	Same
Rosy Apple Aphids	1/4 gal	pint/100 water	For First Brood
Codling Moths	apply at petal fall and every 10-14 days thereafter until control is achieved		
Leafminers			

Green Peach Aphids	1/4 gal	pint/100 water	Same
Rosy Apple Aphids	1/4 gal	pint/100 water	For early delayed dormant application only
Fruitree Leafrollers	2 gal	of heavy 70 sec oil with the spray mixture	For the Western states combine 1 1/2 gal of heavy 140 sec oil with the spray mixture. Do not combine phosphamidon with captan or sulphur
European Red Mites			
San Jose Scales			

Grapefruit	Aphids, Thrips	1 1/2 pint	100 gal water or 1 1/4 2 1/2 pints/acre	For ground application only
Lemons				
Oranges				

CROP  
Grapefruit  
Lemons  
Oranges

CROP  
Broccoli  
Cauliflower

CROP  
Cantaloupe

CROP  
Cotton

CROP  
S

CROP  
A

CROP  
Pe

CROP  
Cucumbers

CROP  
Peppers

CROP  
E

CROP  
Potatoes  
Le  
m

CROP  
P

CROP  
Tomatoes  
Le  
W

NET CONTENTS



**DANGER: KEEP OUT OF REACH OF CHILDREN  
SEE WARNING AND ANTIDOTE STATEMENTS ON THIS LABEL**

...RE USING!

and  
ORS

EMICAL  
ANY

PHONE 602 244-1311

26-431A  
26-AZ-1  
DON 8

74.5%  
3.5%  
22.0%  
100.0%

icide for the control of both  
ed as a diluted spray and is

**READ LABEL**

**DANGER**

**DIRECTIONS FOR USE**

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep pesticide in original container. Do not use food or drink containers for diluting.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:  
GROUND APPLICATION — Vegetable and Field Crops — 30 to 250 gals. Deciduous Fruit Crops — 60 to 800 gals. Citrus — 400 to 3,000 gals. (On young Fruit Plantings — Use a minimum of 15 gals. water per acre).

Where Recommended for AIR APPLICATION — Field Crops — 3 to 10 gals. Orchard Crops — 10 to 20 gals.

NOTE: Important — Avoid drift of this product to crops for which it is not registered or to any crop for which its use may be registered but application would be closer to harvest than recommended. This product is to be applied as a foliar application only.

CROP	CROP PEST	DOSAGE RATE/USE	LIMITATIONS
Apples	Aphids	1/6 pint/100 gal water	Do not apply within 30 days before harvest. Do not apply more than 1 quart of this product per acre.
	Green Peach Aphids	1/4 pint/100 gal water	Same
	Rosy Apple Aphids	gal water	For
	First Brood Codling Moths	Codling Moths	apply at petal fall and every 10-14 days thereafter until control is achieved.
	Leafminers	Leafhoppers	
	Green Peach Aphids	1/4 pint/100 gal water	Same
	Rosy Apple Aphids	gal water	For
	Fruitree Leafrollers	early delayed dormant ap	plication only
	European Red Mites	For areas other than Western states, combine 2 gal of heavy 70 sec oil with the spray mixture. For the Western states, combine 1 1/2 gal of heavy 140 sec oil with the spray mixture. Do not combine phosphamidon with captan folpet or sulphur.	
Grapefruit	Aphids, Thrips	1/4 1/2 pint/100 gal water or 1 1/4 2 1/2 pints/acre	Do not apply within 15 days before harvest. Do not apply more than 2 1/2 pint of this product per acre.
Lemons			
Oranges			

CROP	CROP PEST	DOSAGE RATE/USE	LIMITATIONS
Grapefruit, Lemons, Oranges	Aphids, Thrips	Apply during periods of attack on new flush growth	Same
		1 2 3/4 pint/acre	Same
		For aerial application only	
		Apply during periods of attack on new flush growth	
Broccoli	Aphids, Imported Cabbageworms	1/2 1 pint	Do not apply within 3 days before harvest
Cauliflower			
Cantaloups	Flea-hoppers, Spider Mites, Thrips	1/8 1/2 pint	Do not apply within 24 hours before harvest
	Aphids, Leafminers	1/8 1/2 pint	Same
Cotton	Flea-hoppers	3/16 pint	Use restricted to areas other than Oklahoma and Texas. For early season control only (2 10 leaf stage).
	Spider Mites, Thrips	3/16 pint	For early season control only (2 10 leaf stage).
	Aphids	3/16-1/4 pint	Use lower dosage for early season control (2 10 leaf stage) and higher dosage for mid and late season control.
	Flea Beetles, Leafhoppers, Lygus Bugs	1/2 pint	For mid to late season control only.
Cucumbers	Aphids, Leafhoppers, Leafminers	1/4 1/2 pint	Do not apply within 3 days before harvest.
Peppers	Aphids, Leafminers, Whiteflies	1/2 pint	Do not apply within 6 days before harvest
	European Corn Borers	1 pint	Same
Potatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Leafminers, Stink Bugs	1/2 pint	Do not apply within 7 days before harvest
	Potato Tuberworms	1/2 1 pint	Same
Tomatoes	Aphids, Colorado Potato Beetles, Flea Beetles, Leafminers, Stink Bugs, Whiteflies	1/2 pint	Do not apply within 10 days before harvest

ACCEPTED  
 1526-431

CROP	CROP PEST	DOSAGE RATE/ACRE	LIMITATIONS
Watermelons	Aphids, Leafhoppers, Leafminers	1/4 1/2 pint	Do not apply within 24 hours before harvest
Tangerines	Aphids, Thrips	1-4 1-2 pint/100 gal water or 1-4 2 1-2 pints/acre	Do not apply within 15 days before harvest. Do not apply more than 2 1-2 pints of this product per acre.
		pints/acre	Use restricted to Arizona and California only.
			For ground application only.
			Apply during periods of attack on new flush growth.
		1/2 3/4 pint/acre	Same
			Use restricted to Arizona and California only.
			For aerial application only.
			Apply during periods of attack on new flush growth.
Walnuts	Aphids, Codling Moths	1 pint/100 gal water/acre	Do not apply within 7 days before harvest. Do not apply more than 1 pint of this product per acre.
			For ground application only.
			For Codling Moths, apply at petal fall and every 10-14 days thereafter until control is achieved.
		1 pint/10 20 gal water/acre	Same
			For aerial application only.
			For Codling Moths, apply at petal fall and every 10-14 days thereafter until control is achieved.
	Walnut Husk Flies	1 pint/200 300 gal water/acre	Same
			For ground application only.
			Apply after first egg "stings" appear on husks.
		1 pint/10 20 gal water/acre	Same
			For aerial application only.
			Apply after first eggs "stings" appear on husks.

**NOTICE**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**OUT OF REACH OF CHILDREN**  
**ANTIDOTE STATEMENTS ON THIS LABEL**