



CAUTION: Harmful if swallowed. Do Not Inhale. Keep Out of Reach of Children.

WARNING

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on clothing, or on skin. Do not breathe the dust or fumes. Do not take internally. If spilled on skin or clothing, wash thoroughly with soap and water and change clothes. Bathe promptly with soap and water after working with this product and before eating or smoking. Wash clothing with soap and water before re-use. When container is empty, burn and bury with trash.

WARNING

Keep out of reach of children and domestic animals. Do not contaminate feed or foodstuffs. Use only for recommended purposes at recommended dosages. Toxic to fish and wildlife. Keep out of lakes, streams, or ponds with this material. Burn or destroy containers; never reuse. Stay out of smoke or fumes. Do not apply DYLON to, or allow dust to drift onto, varieties of sorghum which are sensitive to phosphates. To do so may cause burning of the roots or hills. DO NOT APPLY THIS MATERIAL WITH TOPICAL DDT'S EVER BEEN USED TO APPLY 2,4-D, 2,4,5-T OR OTHER RELATED WEED-KILLING COMPOUNDS.



PHOSPHO KIL-5
TOXAPHENE 10

(Contains 5% DYLON*)

U. S. P. A. R. No. 1526


ACTIVE INGREDIENTS:

Diethyl 2,2,2-trichloro-1-hydroxyethyl-phosphonate***	5.00%
Toxophene***	10.00%
INERT INGREDIENTS:	85.00%
	100.00%

*Reg. U.S. & Canadian Pat. Offs. by Farbenfabriken
Baker, A. G. Chemagro Corporation Licensee.

**U.S. Patent No. 2,701,225 Canadian Patent No.
529,379, Effective for Purpose Claimed.

***Technical Chlorinated Carbon (Chlorine
content 67 - 69%)



Main Office: 734 East S. P. Drive - Phoenix, Arizona 85071 - P

This insecticide is for the control of beet worm, cotton leaf perforator, and lygus cotton. For control of beet armyworm and bugs apply at a rate of 20 to 40 pounds in a manner to give thorough coverage of affected portions of the crop plants. For control of cotton leaf perforator apply at a rate of 40 pounds per acre in a manner to give thorough coverage of the affected portion of the crop. Begin application when a damageable infestation is discovered, repeating at weekly intervals until the pest is controlled. Cotton receiving late application of this insecticide must not be grazed by domestic animals and animals being finished for slaughter. Do not feed treated cotton trash to dairy animals. Do not apply within 7 days of harvest.

ANTIDOTE AND FIRST AID

Give Inactive: CALL PHYSICIAN. Take one 1/100 grain atropine tablets and repeat if necessary.

NOTICE: While this material is on this list it is not to be used for the control of insects on cotton. It is not to be used for the control of insects on other crops. It is not to be used for the control of insects on other crops. It is not to be used for the control of insects on other crops. It is not to be used for the control of insects on other crops.

Red

Red

Red

Red


Red

Keep Out of Reach of Children.

inhaled or absorbed in eyes or on clothing, or dust or fumes. Do not get on skin or clothing, wash with water and change soap and water after use. Before eating or drinking, wash with soap and water before eating or drinking. Burn or bury with

and domestic animals. Do not use on foodstuffs. Use only in recommended dosages. Keep out of lakes, streams, or other water material. Burn or bury if not used. Stay out of areas treated with DYLOX to, or allow of sorghum which are to so may cause burning. DO NOT APPLY THIS MATERIAL TO ANY OTHER PLANT OR MATERIAL USED TO APPLY TO WEED-KILLING COM-

RECEIVED
AUG 31 1957
1526-435



PHOSPHO KIL-5
TOXAPHENE 10

(Contains 5% DYLOX*)

U.S. D.A. Reg. No. 1520 -

ACTIVE INGREDIENTS:


Diethyl 2,2,2-trichloro-1-hydroxyethyl-phosphonate**----- 5.00%
Toxophene***----- 10.00%

INERT INGREDIENTS:----- 85.00%
100.00%

*Reg. U.S. & Canadian Pat. Offs. by Farbenfabriken
Baker, A. G. Chemagro Corporation Licensee.

**U.S. Patent No. 2,701,225 Canadian Patent No.
529,379, Effective for Purpose Claimed.

***Technical Chlorinated Camphene (Chlorine
content 67 - 69%)



Main Office: 734 East S. P. Drive - Phoenix, Arizona 85021 - Phone 253-1111

This insecticide is for the control of beet armyworm, cotton leaf perforator, and lygus bugs on cotton. For control of beet armyworm and lygus bugs apply at a rate of 20 to 40 pounds per acre in a manner to give thorough coverage of the affected portions of the crop plants. For control of cotton leaf perforator apply at a rate of 30 to 40 pounds per acre in a manner to give thorough coverage of the affected portion of the crop plants. Begin application when a damageable infestation is discovered, repeating at weekly intervals until controlled. Cotton receiving late application of this insecticide must not be grazed by dairy animals and animals being finished for slaughter. Do not feed treated cotton trash to dairy or other animals. Do not apply within 7 days of harvest.

ANTIDOTE AND FIRST AID

Give Inactive: CALL PHYSICIAN. Take one or two 1/100 grain atropine tablets and repeat in 1 hour if necessary.

NOTICE

While statements made on this label are true to the best of our knowledge, Chemagro Corporation does not warrant, either expressly or implied, that the use of this material will result in the control of the pest described in the label. The user assumes all responsibility for the results of the use of this material. The user agrees to hold Chemagro Corporation harmless for any and all claims, damages, or losses resulting from the use of this material, whether or not in connection with the use of this material. The user agrees to hold Chemagro Corporation harmless for any and all claims, damages, or losses resulting from the use of this material, whether or not in connection with the use of this material. The user agrees to hold Chemagro Corporation harmless for any and all claims, damages, or losses resulting from the use of this material, whether or not in connection with the use of this material.

Red
Red
Red
Red

Red

50 LBS.