

CAUTION: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale.

WARNING

HARMFUL IF SWALLOWED! MAY BE ABSORBED THROUGH SKIN! May produce irritation of skin or eyes. Do not breathe spray mist. Avoid contact with skin and eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

ANTIDOTE

Remove from further contact. Do not induce vomiting. CALL A PHYSICIAN.

CAUTION

FLAMMABLE MIXTURE. Keep away from heat or open flame. DO NOT use on vegetable or fruit crops. Avoid contamination of soils which subsequently may be planted with root crops (such as potatoes and carrots) since residues of this material may cause off flavor.


CAUTION

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated.

CAUTION

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

Camera man's Note  
Poor Copy




U. S. D. A. Reg. No. 1526

This is an emulsifiable concentrate containing one pound of gamma isomer of BENZENE HEXACHLORIDE per gallon at 60° F.

ACTIVE INGREDIENTS:	
Gamma isomer of benzene hexachloride-----	10.00%
Other isomers of benzene hexachloride----	17.50%
Aromatic Petroleum Solvent-----	68.50%
INERT INGREDIENTS:-----	
	4.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN



Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

This insecticide is for the control of cotton leaf worms, stink bugs, cotton stainer, lygus, thrips, leafhoppers, grasshoppers, and certain aphids on cotton. For control of cotton leafworms, cotton stainer, lygus, leafhoppers, grasshoppers and aphids, apply at the rate of 1 1/2 to 2 quarts per acre. For control of stink bugs apply at the rate of 2 quarts per acre. For control of thrips apply at the rate of 1 quart per acre. For ground application dilute in 10 to 30 gallons of water. For air application dilute in 1 to 10 gallons of water. Repeat at weekly intervals as long as necessary for control. Do not apply in rotation with root crops. Do not graze treated fields to dairy or meat animals. For more specific information, consult your state agricultural extension service.

FLY CONTROL - For an effective residue on farm buildings, add 2 gallons BHC EC-10 to 100 gallons of water and apply to wet all surfaces to point of run-off. Before applying spray, cover feed and water troughs. Protect all feed from spray. Do not use on farm animals or in or around poultry houses. Do not use in or around dairy barns. Use on exterior surfaces only of farm buildings.

**ACCEPTED**  
DEC 5 1967  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 1526-14/SUBJECT  
TO ATTACHED COMMENTS.

# 1526-441

Seller makes no warranty, expressed or implied, concerning the use of this product other than that stated on the label. Buyer assumes the risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Red

THE MANUFACTURER OF THIS PRODUCT IS NOT RESPONSIBLE FOR ANY DAMAGE TO PERSONS OR PROPERTY CAUSED BY THE USE OF THIS PRODUCT.

CONTENTS  
GALS.