

PRECAUTIONARY STATEMENTS

CAUTION 1526-443

HAZARDS TO HUMANS: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily.

ANTIDOTE: Note to Physician: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS: To protect fish and wildlife, do not contaminate any body of water with this material. This pesticide is toxic to aquatic organisms. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated area.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM THE AREA BEING TREATED.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL:

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. OPEN DUMPING IS PROHIBITED.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures such as limited controlled burning.

Apply when pests first appear and repeat as necessary. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop. Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

ALMONDS (0): Fruit-tree Leafroller. Use 40 to 80 pounds per acre.

ALFALFA and CLOVERS (0): Use 15 lbs. per acre to control Thrips, Leafhoppers, Three Corned Alfalfa Hoppers, Japanese Beetles, Alfalfa Caterpillars, Armyworms, Corn Earworms, and Stink Bugs.

APPLES (1), PEARS (1): Codling Moth, Green Apple Aphids, Apple Maggot, Redbanded Leafroller, Pear Psylla, Periodical Cicada. Use 35 to 60 lbs. per acre. Application within 30 days after full bloom may also provide apple thinning. To avoid this, delay application until at least 30 days after bloom.

APRICOTS (3), PEACHES (1), NECTARINES (3): Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth. Use 40 to 60 lbs. per acre.

ASPARAGUS (1): Asparagus Beetle on Seedlings or Spears. Use 20 pounds per acre. Apply during cutting season. Do not repeat application within 14 days. Asparagus Beetle, Asparagus Cicada on ferns or brush growth in the post-harvest period. Use 20 to 40 pounds per acre.

BEANS (0): Bean Leaf Beetle, Cucumber Beetle, Flea Beetles, Leafhoppers, Corn Earworm, Stink Bugs, Tarnished Plant Bug. Use 20 pounds per acre. California Legume Bug, Pink Lima Bean Pod Borer. Use 20 pounds per acre. Do not feed treated beans.


BLACKBERRIES (7), BOYSBERRIES (7), RASPBERRIES (7), LOGANBERRIES (7), RASPBERRIES (7): Leafrollers, including Omnivorous Leafroller, Sawfly, Climbing Cutworm, Snowy Tree Cricket (Calif.), Orange Tortrix (Oregon), Leafhopper, Raspberry Aphids. Use 20 pounds per acre.

STOP - READ LABEL BEFORE USING!

Agro-Chem Brand

Mfg. for CHEMICAL DISTRIBUTORS, INC.

DBA



1537 PHOENIX, ARIZ. 85036 PHONE 602 243-2711

E.P.A. REG. NO. 1526-443
E.P.A. EST. NO. 33691-AZ-1

SEVIN* 10 DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl methylcarbamate) 10.00%

INERT INGREDIENTS:

90.00%

100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED OR INHALED induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN, flush eyes thoroughly with water, wash skin thoroughly with soap and water.

* Trademark of Union Carbide Corporation for the active ingredient carbaryl.

NET CONTENTS

50 LBS.

CAUTION: Keep Out of Reach of Children
See other precautions on this label.

STOP - READ LABEL BEFORE USING!

Agro-Chem Brand

Mfg. for CHEMICAL DISTRIBUTORS, INC.

DBA



ARIZONA AGROCHEMICAL COMPANY

Growing thru Chemical Progress

E.P.T.B.D. 1537 PHOENIX, ARIZ. 85036 PHONE 602 243-2711

E.P.A. REG. NO. 1526-443

E.P.A. EST. NO. 33691-AZ-1

24 10/11

Do not apply to plants or animals that are susceptible to the active ingredient.

1526-443

SEVIN* 10 DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl methylcarbamate) 10.00%

INERT INGREDIENTS:

90.00%

100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED OR INHALED induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN, flush eyes thoroughly with water, wash skin thoroughly with soap and water.

* Trademark of Union Carbide Corporation for the active ingredient carbaryl

DIRECTIONS FOR USE

1526-443

CABBAGE (3), BROCCOLI (3), BRUSSELS SPROUTS (3), CAULIFLOWER (3), KOHLRABI (3): Flea Beetles, Harlequin Bugs — Use 10 pounds per acre. Armyworm, Corn Earworm, Cabbage Looper (small) — Use 20 pounds per acre. For small Cabbage Looper control after edible parts start to form, apply in a 5 to 7 day schedule.

CARROTS (0), CHINESE CABBAGE (14), COLLARDS (14), DANDELION (14), ENDIVE (14), BEETS (Roots 3 - Tops 14), HORSERADISH (3), KALE (14), LETTUCE (Head 3 - Leaf 14), MUSTARD GREENS (14), PARSLEY (14), PARSNIPS (3), RADISHES (3), RUTABAGAS (3), SALSIFY (Roots 3 - Tops 14), SPINACH (14), SWISS CHARD (14), TURNIPS (Roots 3 - Tops 14): Flea Beetle, Harlequin Bugs (not on Spinach or Dandelion), Leafhoppers — Use 10 to 15 pounds per acre. Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Cabbage Looper (small) — Use 20 pounds per acre. For small Cabbage Looper control apply in a 5 to 7 day schedule.

CHERRIES (1), PLUMS (1), PRUNES (1): Cherry Fruit Fly, Peach Twig Borer, Leafhopper, Codling Moth, Orange Tortrix, Western Tussock Moth — Use 40 to 60 pounds per acre.

CORN (0): Use 20 lbs. per acre to control Corn Earworm, Fall Armyworm, Sap Beetles and Japanese Beetles. Do not apply within 7 days of harvest for forage. For Corn Earworms apply to silks as soon as they appear. Make 3 to 4 applications of 2-3 day intervals. Reapply immediately after rain.

COTTON (0): Use 20 to 25 pounds per acre to control. Thrips, Flea Beetles, Flea Hoppers, Leafworms, Bollworms, Fall Armyworms, Leaf Perforators, Leaf Rollers, Leafhoppers, Lygus Bugs, Salt Marsh Caterpillars, Stink Bugs and Grasshoppers. Do not graze treated fields or feed cotton trash to dairy or meat animals.

FILBERTS (0): Filbert Aphid, Filbert Leafroller, Filbert Moth — Use 40 to 50 pounds per acre.

GRAPES (0): Use 20 to 25 pounds per acre to control Grape Leaf Folders, Grape Leafhoppers, Grape Berry Moths Japanese Beetles and Western Grape Leaf Skeletonizer.

EGGPLANTS, PEPPERS and TOMATOES Use 20 pounds per acre to control Tomato Fruitworms, Fall Armyworms, Flea Beetles, Leafhoppers, Tomato Hornworms, European Corn Borer, Stink Bugs, Colorado Potato Beetles and Lygus Bugs. No time limitation.

OKRA (0): Corn Earworm, Stink Bugs — Use 20 pounds per acre

POTATOES: Use 15 to 20 pounds per acre to control Leafhoppers, Flea Beetles and Colorado Potato Beetles. Apply to foliage only.

RICE (14): Armyworms — Use 10 to 15 pounds per acre Tadpole Shrimp (Calif) — Use 20 pounds per acre Do not apply within 14 days before or after application of the herbicide propanil (Stam F 34 or Roquel) as injury may result

SORGHUM (MILO, GRAIN SORGHUM) (Forage 0 - Grain - 21) Armyworm, Corn Earworm, Stink Bugs, Webworm — Use 20 pounds per acre Sorghum Midge — Use 15 pounds per acre Treatment for Sorghum Midge control should be made 3 to 4 days after heads have emerged from boot. Direct treatment into the forming heads

STRAWBERRIES (1) Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil — Use 20 pounds per acre

SUGAR BEETS (14) Armyworms, Beet Leafhoppers, Beet Webworms — Use 20 pounds per acre Grasshoppers — Use 15 pounds per acre

TOMATO EGGPLANT, PEPPERS (0) Colorado Potato Beetle, Flea Beetles, Leafhoppers — Use 5 to 10 pounds per acre Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs, Stink Bugs — Use 10 to 20 pounds per acre

CAUTION

Use only for recommended purposes and at recommended dosages

Do not apply this material with equipment that has ever been used to apply 2,4-D, 2,4,5-T or related weed killing compounds

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and of handling of this material when such use and/or handling is contrary to label instructions.

ON: Keep Out of Reach of Children.

See other precautions on this label.