

Keep out of reach of children. Harmful if swallowed. Do not inhale.

WARNING

Harmful if swallowed, inhaled, or absorbed through the skin. May cause irritation of skin and eyes. Do not get in eyes, on skin, or on clothing. Do not inhale fumes. Do not take internally. Use clean clothing daily. Wash thoroughly with soap and water after handling this product and before eating or smoking. Keep out of reach of children and domestic animals. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate feed or foodstuffs.

ANTIDOTE

In case of skin contact, wash thoroughly with soap and water. For eyes, flush with plenty of water and get medical attention. If swallowed, do not induce vomiting. CALL A PHYSICIAN.


CAUTION

This product is toxic to aquatic invertebrates. Keep out of streams, lakes or ponds. Shrimp and crab may be killed at application rates recommended at this level. Do not apply where these are important resources.

Use only for recommended purposes at recommended dosages. Do not apply this material with equipment that has ever been used to apply 2,4-D, 2,4,5-T or related weed killing compounds.

CAUTION

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place. Do not contaminate water by cleaning of equipment, or disposal of wastes.




U. S. D. A. Reg. No. 1526—

Tedion (R) EC-12 is an emulsifiable miticide containing 1 pound actual Tedion per gallon at 60° F. A water stable emulsion of desired strength can be prepared easily and can be used in conventional hydraulic sprayers, low volume ground applicators, and airplane sprayers. For airplane application dilute in 3 to 10 gallons of water per acre. For ground application dilute in 10 to 30 gallons of water per acre.

ACTIVE INGREDIENTS:

Tetradifon (4-chlorophenyl 2,4,5-tetra-	
chlorodiphenyl sulfone)-----	12.3%
Aromatic Petroleum Solvent-----	35.0%
<u>INERT INGREDIENTS:-----</u>	<u>52.7%</u>
	100.0%



Main Office: 224 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

<u>Crop</u>	<u>Mite</u>	<u>Rate of Application</u>	<u>Limitations</u>
<u>Citrus</u>	Citrus red mite Six-spotted mite Texas citrus mite	2 to 5 gals. Tedion EC-12 per acre	Do not apply more than once per season while fruit is present.
<u>Cotton</u>	Spider mite	1/2 to 1 gal. Tedion EC-12 per acre	Do not apply after bolls open. Do not feed or graze treated foliage or trash to livestock.
<u>Seed</u> (Alfalfa)	Spider mite	1/2 to 3/4 gal. Tedion EC-12 per acre	Do not feed treated forage, straw or threshings to livestock.

This spray should be applied in any convenient amount of water that will give thorough coverage of the affected portions of the crop plants, and adhere to the rate of applications listed in the above table. Begin applications when the mites first appear, and continue applications as additional control is needed, heeding the limitations listed in the above table. For further specific information, consult your state agricultural experiment station.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all liability for any damage to crops or other property resulting from the use of this product.

red

GM 1