

114

May produce irritation of skin or eyes. Do not
use on clothing. Wash thoroughly with soap and
water and remove clothing. Flush with water and get
medical attention. Avoid contamination of food and
feedstuffs. Do not use on household pets or humans.

CAUTION

Remove from further contact. Do not induce vomiting.

CAUTION

Do not use on clothing. Wash thoroughly with soap and
water and remove clothing. Flush with water and get
medical attention. Avoid contamination of food and
feedstuffs. Do not use on household pets or humans.

CAUTION

This is an emulsifiable concentrate containing one
pound of gamma isomer of BENZENE HEXACHLORIDE per
gallon at 60° F.

ACTIVE INGREDIENTS:	
Gamma isomer of benzene hexachloride-----	10.00%
Other isomers of benzene hexachloride-----	17.50%
Aromatic Petroleum Solvent-----	68.50%
INERT INGREDIENTS:-----	
	100.00%

This insecticide is for the control of cotton leaf
worms, stink bugs, cotton stainer, lygus, thrips,
leafhoppers, grasshoppers, and certain aphids on
cotton. For control of cotton leafworms, cotton
stainer, lygus, leafhoppers, grasshoppers and
aphids, apply at the rate of 1 to 2 quarts per
acre. For control of stink bugs apply at the rate
of 2 quarts per acre. For control of thrips apply
at the rate of 1 quart per acre. For ground appli-
cation dilute in 10 to 50 gallons of water. For
air application dilute in 1 to 10 gallons of water.
Apply at regular intervals during the growing season.
Do not use on clothing. Wash thoroughly with soap
and water and remove clothing. Flush with water and
get medical attention. Avoid contamination of food
and feedstuffs. Do not use on household pets or
humans.

CAUTION

ACCEPTED
DE 15 50
U.S. DEPARTMENT OF AGRICULTURE
FOR THE ENFORCEMENT OF THE
FEDERAL INSECTICIDE ACT
BY THE UNITED STATES DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C.

1526-457

DANGER: Keep Out of Reach of Children. Harmful if Swallowed. Do Not Inhale.

WARNING

May be fatal if swallowed, inhaled or absorbed thru the skin. Parathion is rapidly absorbed thru the skin! Do not get in eyes, nor allow inhalation or skin contact. Wear protective clothing, goggles, and natural rubber gloves. Wear a mask or respirator of the type passed by the U.S. Department of Agriculture for protection against Parathion. Wash hands, arms, and face thoroughly with soap and water after exposure, and before eating and smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and food-stuffs. Do not feed refuse from the treated crops to meat and dairy animals.

ANTIDOTE


FIRST AID TREATMENT: Atropine sulfate tablets grains 1/100. For severe Parathion poisoning, take two tablets at once. Avoid further exposure to Parathion type insecticides until permitted by doctor. **CALL A PHYSICIAN AT ONCE.**

CAUTION

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds.

CAUTION

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.




U. S. D. A. Reg. No. 1526—

ACTIVE INGREDIENTS:

Parathion*-----	2.00%
Diethyl diphenyl dichloroethane-----	4.75%
Related compounds-----	.25%
INERT INGREDIENTS:-----	93.00%
	100.00%

*O,O-diethyl O-O-nitrophenyl phosphorothioate



Main Office: 734 East S. P. Drive — Phoenix, Arizona 85001 — Phone 253-1111

DIRECTIONS FOR USE

This product is for the control of leafhoppers, cabbage looper, aphids, lygus and thrips on lettuce and for the control of cabbage looper and aphids on broccoli, cabbage and cauliflower. Apply in a manner to give thorough coverage to the affected portions of the crop-plants as soon as damageable infestation is discovered, and repeat at weekly intervals if additional control is needed. For further information as to application, consult your state agricultural extension service.

CROP	INSECT PEST	MAXIMUM RATE	LIMITATIONS
Broccoli Cauliflower	Cabbage looper Aphids	25 lbs. per acre	Do not apply within 7 days of harvest.
Lettuce (Leaf or Bibb)	Leafhoppers Cabbage looper Aphids Lygus Thrips	25 lbs. per acre	Do not apply within 21 days of harvest.
Cabbage	Cabbage looper Aphids	25 lbs. per acre	Do not apply within 10 days of harvest.
Lettuce (Head)	Leafhoppers Aphids Cabbage looper Lygus Thrips	25 lbs. per acre	Do not apply within 7 days of harvest.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



DANGER POISON



DANGER KEEP OUT OF REACH OF CHILDREN. SEE WARNING AND ANTIDOTE STATEMENTS ON SIDE PANEL

NET CONTENTS

LBS.