

**MALATHION - FOR INSECT CONTROL**  
**5% DUST**

Area	Insect	Application
Homes, Dairies, Food Processing Plants	Ants, crickets, clover mites, earwigs, Pantry Pests,* scorpions, silverfish, spiders. (*Pantry pests such as the exposed stages of: saw-toothed grain beetle, flour beetle, rice weevil, cigarette beetle, drug store beetle, and Indian meal moth.)	Apply malathion 5% dust liberally to window sills, cracks and crevices, around doors, baseboards, storage areas, bookcases, under sinks and to other areas frequented by these pests. Avoid contamination of food, utensils, milk, milk equipment, and water. <b>CENTIPEDES, SCORPIONS:</b> Apply malathion 5% dust liberally to window sills, cracks and crevices, around doors, baseboards, storage areas, bookcases, under sinks and to other areas frequented by these pests.

**HOMES, DAIRIES, FOOD PROCESSING PLANTS**

Insect	Application
Fleas, brown dog ticks	Apply malathion 5% dust liberally to floors, cracks, and crevices, sleeping quarters of animals, and to other areas frequented by these pests.
Carpet beetles, Clothes moths.	Apply malathion 5% dust liberally along baseboards, edges of carpeting, under furniture, under carpeting and rugs, in closets and shelving and to other areas frequented by these pests.
Roaches	Apply malathion 5% dust liberally to window sills, cracks and crevices, around doors, baseboards, storage areas, bookcases, under sinks and to other areas frequented by these pests.
Mosquitoes and small flying insects.	<b>OUTDOORS:</b> Apply malathion 5% dust liberally around doors, windows and porches and to other flat surfaces where pests are known to alight or congregate.

20451

Use 5% Malathion Dust for the following insects on Ornamentals:  
Aphids - Spider mites - Bagworms

Greenhouses (in and around Greenhouses, homes and gardens where non food crops are grown)	Insects Millipedes Springtails Sowbugs	Application
		Apply 5% dust at the rate of 1 lb. per 150 square feet of soil surface or where insects congregate. Repeat at 7-10 day intervals as needed.

Animals	Pest	Application
Cattle - Beef and non-milking. Horses	Lice Ticks Hornfly	3 tbsps. 5% Malathion Dust per animal. Apply on back and neck of the animals and repeat applications at 10-14 day intervals.  Do not treat animals under one month of age.  Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.  When applying sprays and dusts avoid contamination of feed, food containers and watering troughs.


**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
**MALATHION 5**

INSECTICIDE DUST

FOR **PCO**

Formulated for and sold exclusively to:

ACS  
1961-12-21  
1526-413  
UNITED STATES  
F. REG.  
FOY 1



**ARIZONA**  
**AGROCHEMICAL**  
**CORPORATION**

*Growing thru Chemical Progress*

Head office: 734 East S. P. Drive • Phoenix, Ariz. 85001 • Phone 253-1111

**CAUTION**

KEEP OUT OF THE REACH OF CHILDREN

# MALATHION 5

INSECTICIDE DUST

FOR **PCO**

Formulated for and sold exclusively to:

ACCS

Dec 21 - 1969

UNDER THE FEDERAL FOOD, DRUG & COSMETIC ACT

ACTIVE INGREDIENTS:	By Weight
Malathion	5.0%
INERT INGREDIENTS	95.0%
	100.0%

U.S.D.A. Registration No. 1526 - 463

\*O, O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate

**WARNING**

Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

**ANTIDOTES**

If swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately!  
If on skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

**CAUTION**

Keep out of reach of children and domestic animals. Do not contaminate feed or foodstuffs. Use only for recommended purposes at recommended dosages.  
To protect fish and wildlife, do not contaminate lakes, streams or ponds with this material.

**MALATHION - FOR INSECT CONTROL  
5% DUST**

**ANIMALS**

Location	Insect	Application
Cattle, dairy	Horn fly	Thorough, full-coverage applications should be made. Repeat applications as necessary. 3 tblsp. 5% Malathion dust per animal. Apply on back and neck of the animals and repeat applications at 10-14 day intervals.  Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.
Dogs & cats	Fleas	5% Dust - complete coverage. Repeat treatments every 2 or 3 weeks, if necessary.
Pet Quarters	Fleas	1-2 lb. per 1000 sq. ft. of surface. Remove debris, then treat and repeat when necessary.
Hogs	Lice	Treat animal thoroughly. Repeat in 10 days and thereafter as needed.  When applying dusts avoid contamination of feed, food containers and watering troughs.

**NOTICE**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

PCO Form 13 2 1/2 M 6-68

**ARIZONA**  
**AGROCHEMICAL**  
**CORPORATION**



*Growing thru Chemical Progress*

Head office: 734 East S. P. Drive • Phoenix, Ariz. 85001 • Phone 253-1111