

PRECAUTIONARY & ANTIDOTE STATEMENT

CAUTION

MAY BE FATAL IF SWALLOWED. This material if brought into contact with the skin may cause toxic symptoms and irritation. Avoid contact with eyes, skin and clothing. Do not inhale dusts. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

FIRST AID FOR FLUORIDES

In case of contact, wash thoroughly with soap and water. For eyes, flush with plenty of water and obtain medical attention promptly. Give at once, large draught of lime water or weak calcium chloride solution, stimulants of strong coffee or aromatic spirits of ammonia and artificial respiration. Keep lower extremities and chest warm. CALL A PHYSICIAN.

CAUTION

Keep all persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the drifting insecticides and volatile components have dissipated.

Do not apply under conditions involving possible drift to food or forage that might be damaged or rendered unfit for sale, use, or consumption.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

SEE CAUTION


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

STOP - READ LABEL BEFORE USING

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

DEA



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526-485

E.P.A. EST. NO. 1526-AZ-1

CRYOLITE 50 DUST


ACTIVE INGREDIENTS:

Sodium Fluoaluminate 50.0%

INERT INGREDIENTS: 50.0%

100.0%

(Fluorine expressed as elemental 27.1%)



DO NOT USE THIS PRODUCT FOR

NOTE: Important - Avoid or to any crop for which harvest than recommended

CAUTION: Do not apply to bush and stone fruits or Grapes after buds or Greens, Cauliflower or Lettuce stage if tops are to of fruit and vegetables by

Apply to the following crops

CROP
Apples, Pears, Peaches, Sh
Beans
Cantaloupes, Watermelons, Squash
Cabbage, Collards, Mustard Turnips, Broccoli, Radishes
Citrus (Grapefruit, Kumquat, Limes, Oranges, Tangelos)
Cranberries
Grapes
Lettuce
Peppers
Strawberries
Tomatoes

CAUTION! Keep Out of Reach of Children
See other precautions on this label


STATEMENT

STOP - READ LABEL BEFORE USING!

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

D.B.A.



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526-485
E.P.A. EST. NO. 1526-AZ-1

CRYOLITE 50 DUST


ACTIVE INGREDIENTS:

Sodium Fluoaluminate 50.0%

INERT INGREDIENTS: 50.0%

100.0%

(Fluorine expressed as elemental 27.1%)



DO NOT USE THIS PRODUCT FOR ANY USES OTHER THAN THOSE SPECIFIED IN THE DIRECTIONS.

NOTE: Important - Avoid dust of this product to crops for which it is not registered or to any crop for which its use may be registered but application will be before harvest than recommended. This product is to be applied as a foliar application.

CAUTION: Do not apply to corn. Do not apply immediately before harvest. Do not apply to bush and stone fruits after fruit begins to form. Do not apply to Broccoli, Cabbage, or Grapes after heads or edible parts start to form. Do not apply to Collards, Mustard Greens, Cauliflower or Lettuce after seedling stage. Do not apply to Turnips after seedling stage if tops are to be used for feed. Remove excess residues from edible parts of fruit and vegetables by washing, brushing, or other effective means.

Apply to the following crops at a rate of 50-100 pounds/acre only.

CROP	CROP PEST
Apples, Pears, Peaches, Shadetrees	Flea weevils, Codling moths, Gypsy moth Caterpillars, Fuller rose beetles, Plum curculio.
Beans	Mexican bean beetles, Bean leafrollers, Cucumber beetles, Corn earworms, Bean leaf beetles.
Cantaloupes, Watermelons, Cucumber Squash	Armyworms, Pickleworms, Cucumber beetles
Cabbage, Collards, Mustard Greens, Turnips, Broccoli, Radishes, Cauliflower	Imported cabbageworms, Cabbage loopers, Diamondback moth caterpillars, Yellow striped armyworms, Climbing cutworms, Corn earworms, Flea beetles.
Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines)	Citrus cutworms, Variagated cutworms, Orangedog, Fruittree leafrollers, Orange tortrix, Garden tortrix, Holcocera, Fuller rose weevils, Katydid.
Cranberries	Cranberries fruitworms, Gypsy moth caterpillars, False armyworms.
Grapes	Grape leafrollers, Omnivorous leafrollers.
Lettuce	Cabbage loopers.
Peppers	Pepper weevils.
Strawberries	Strawberries leafrollers.
Tomatoes	Hornworms, Fruitworms, Blister beetles, Pinworms.

CAUTION! Keep Out of Reach of Children
See other precautions on the label

NOTICE