

STOP - READ LABEL BEFORE USING!

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

OR



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311

E.P.A. REG. NO. 1526- 502
E.P.A. EST. NO. 1526-AZ-1

ZM 8% FUNGICIDE DUST

ACTIVE INGREDIENTS:

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate ion in which the ingredients are:

Manganese	1.6%
Zinc	0.2%
Ethylene bisdithiocarbamate ion	6.2%
INERT INGREDIENTS	92.0%
	100.0%

IMPORTANT

Keep away from fire and sparks. Store in a cool, dry place.

Do not allow to become wet or overheated during storage.

This may bring on chemical changes in this product and may also generate flammable vapors. Keep container closed when not in use.

DIRECTIONS FOR USE

Do not use this product for any uses other than those specified on this label.

CROP	CROP PEST	DOSAGE RATE/ACRE	LIMITATIONS
Grapes (California Only)	Bunch rot, dead-arm, early season downy mildew	15 - 40 lbs. Begin application when shoots are 0.5-1.5 inches long. Repeat when shoots are 3-5 inches long, when shoots are 8-10 inches long, and at 10-14 day intervals thereafter until fruit is set.	Do not apply after fruit is set.
Grapes (Areas other than California)	Blackrot, dead-arm, downy mildew	15 - 40 lbs. Begin application when shoots are 0.5-1.5 inches long, repeat when shoots are 3-5 inches long, when shoots are 8-10 inches long and repeat at 10-14 day intervals thereafter.	Do not apply within 66 days before harvest.
Cucumbers, Summer squash, Melons (Cantaloupes, Cassabas, Crenshaws, Honeydews, Honey Balls, Muskmelons, Persians, Watermelons)	Alternaria leaf spot, anthracnose, cercospora leaf spot, downy mildew, gummy stem blight, scab	15 - 27 lbs. Begin application at 2 leaf stage and repeat at 5-7 day intervals.	Do not apply within 5 days before harvest.
Potatoes	Early and late blights	10 - 20 lbs. Begin application when plants are 4-6 inches high, and repeat at 7-10 day intervals or more frequently under severe disease conditions.	No time limitations.
Potato Seed pieces	Scab, seed piece decay	1/2 - 1 lb. 100 lbs. of whole or cut seed pieces. Apply for thorough coverage, and plant as soon after treatment as possible.	Do not use treated seed pieces for food or feed.

SEED TREATMENT USES

For control of seed and soil borne fungi causing seed decay, seedling blights, covered smuts of barley, rye and wheat, seed borne rust (Puccinia Carthami) of safflower, and surface borne bacteria and fungi on tomato seed.

CAUTION: Do not use treated seed for food, feed or oil purposes.

CROP	SLURRY TREATMENT DOSAGE RATE	DRY TREATMENT DOSAGE RATE
Barley	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Field Corn	15-16 - 1 1/2 lbs./bushel	15-16 - 1 1/2 lbs./bushel
Cotton (acid delinted)	1 1/2 lbs./bushel	1 1/2 lbs./bushel
Cotton (reginned)	3 3/4 lbs./bushel	3 3/4 lbs./bushel
Flax	2 1/2 lbs./bushel	1 1/2 - 2 1/2 lbs./bushel
Oats	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Shelled peanuts	1 1/4 - 2 1/2 lbs./bushel	1 1/4 - 2 1/2 lbs./bushel
Rice	1 1/4 - 2 1/2 lbs./100 lbs. seed	1 1/4 - 2 1/2 lbs./100 lbs. seed
Rye	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Safflower	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Sorghum	15/16 - 1 1/2 lbs./bushel	15/16 - 1 1/2 lbs./bushel
Tomatoes	5 lbs./100 lbs. seed	None
Wheat	1 1/4 lbs./bushel	1 1/4 lbs./bushel

NOTICE


Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CAUTION

May cause irritation of nose, throat, eyes, and skin. Do not breath dust. In case of skin contact wash with water. For eyes wash with plenty of water and get medical attention. Do not reuse empty containers. Destroy by burying or burning staying away from smoke and fumes.

Contamination of fish bearing waters with this product may harm fish.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.



NET CONTENTS **50 LBS.**

CAUTION: Keep Out of Reach of Children.
See other precautions on this label.

READ LABEL BEFORE USING!

CHEMICAL DISTRIBUTORS

D B A



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602 244-1311


EPA REG NO 1526-502
EPA EST NO 1526-AZ-1

CAUTION

May cause irritation of nose, throat, eyes, and skin. Do not breath dust. In case of skin contact wash with water. For eyes wash with plenty of water and get medical attention. Do not reuse empty containers. Destroy by burying or burning staying away from smoke and fumes.

Contamination of fish bearing waters with this product may harm fish.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.



ZM 8% FUNGICIDE DUST

ACTIVE INGREDIENTS:

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate 8% in which the ingredients are:

Manganese	1.6%
Zinc	0.2%
Ethylene bisdithiocarbamate ion	6.2%
INERT INGREDIENTS	92.0%
	100.0%

IMPORTANT

Keep away from fire and sparks. Store in a cool, dry place.

Do not allow to become wet or overheated during storage.

This may bring on chemical changes in this product and may also generate amable vapors. Keep container closed when not in use.

Do not use this product for any uses other than those specified on this label.

CROP	CROP PEST	DOSAGE RATE ACRE	LIMITATIONS
Grapes (California Only)	Leaf roll, dead arm every season downy mildew	15-20 lbs. Begin application when shoots are 1/2 grown. Repeat when shoots are 3/4 grown. When shoots are 1/2 grown, spray at 10-15 day intervals thereafter until fruit set.	Do not apply within 14 days of harvest.
Grapes Ariz. other than California	Leaf roll, dead arm downy mildew	15-20 lbs. Begin application when shoots are 1/2 grown. Repeat when shoots are 3/4 grown. When shoots are 1/2 grown, spray at 10-15 day intervals thereafter until fruit set.	Do not apply within 14 days of harvest.
Cucumbers Summer squash, Melons Cantaloupes, Cassabas, Crenshaws Honeydews Honey Balls, Muskmelons Persians Watermelons	Alternaria leaf spot in this case mildew leaf spot downy mildew, gummy stem blight, scab	15-20 lbs. Begin application when shoots are 1/2 grown. Repeat when shoots are 3/4 grown. When shoots are 1/2 grown, spray at 10-15 day intervals thereafter until fruit set.	Do not apply within 14 days of harvest.
Potatoes	Early and late blight	15-20 lbs. Begin application when shoots are 1/2 grown. Repeat when shoots are 3/4 grown. When shoots are 1/2 grown, spray at 10-15 day intervals thereafter until fruit set.	Do not apply within 14 days of harvest.
Potato Seed pieces	Scab, seed piece rot	15-20 lbs. Begin application when shoots are 1/2 grown. Repeat when shoots are 3/4 grown. When shoots are 1/2 grown, spray at 10-15 day intervals thereafter until fruit set.	Do not apply within 14 days of harvest.

SEED TREATMENT USES

For control of seed and soil-borne fungi causing seed decay, damping off, and other seedling diseases in corn, soybeans, wheat, and wheat seedlings. For control of seedling diseases in corn, soybeans, wheat, and wheat seedlings.

CAUTION: Do not use treated seed for food, feed, or other purposes.

CROP	SLURRY TREATMENT DOSAGE RATE	DRY TREATMENT DOSAGE RATE
Barley	15-20 lbs. bushel	15-20 lbs. bushel
Bird Corn	15-20 lbs. bushel	15-20 lbs. bushel
Cotton	15-20 lbs. bushel	15-20 lbs. bushel
Cotton and de-linted	15-20 lbs. bushel	15-20 lbs. bushel
Cotton (regional)	15-20 lbs. bushel	15-20 lbs. bushel
Flax	15-20 lbs. bushel	15-20 lbs. bushel
Guts	15-20 lbs. bushel	15-20 lbs. bushel
Shelled peanuts	15-20 lbs. bushel	15-20 lbs. bushel
Rice	15-20 lbs. bushel	15-20 lbs. bushel
Rye	15-20 lbs. bushel	15-20 lbs. bushel
Safflower	15-20 lbs. bushel	15-20 lbs. bushel
Sorghum	15-20 lbs. bushel	15-20 lbs. bushel
Tomatoes	15-20 lbs. bushel	15-20 lbs. bushel
Wheat	15-20 lbs. bushel	15-20 lbs. bushel

NET CONTENTS 50 LBS.

CAUTION: Keep out of reach of children.
See other precautions on this label.