

STOP - READ LABEL BEFORE USING

Agro-Chem Brand

CHEMICAL DISTRIBUTORS

DBA



P.O. BOX 21537 PHOENIX, ARIZ. 85036 PHONE 602/243-2711

E.P.A. REG. NO. 1526-502
E.P.A. EST. NO. 1526-224

ZM 8% FUNGICIDE DUST

ACTIVE INGREDIENTS:

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate 8% in which the ingredients are:

Manganese	1.6%
Zinc	0.2%
Ethylene bisdithiocarbamate ion	6.2%
INERT INGREDIENTS	92.0%
	100.0%

IMPORTANT

Keep away from fire and sparks. Store in a cool, dry place.

Do not allow to become wet or overheated during storage.

This may bring on chemical changes in this product and may also generate flammable vapors. Keep container closed when not in use.

CAUTION: Keep Out of Reach of Children
See other precautions on this label.

ACCEPTED

JAN 19 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act

1526-502

CAUTION

KEEP OUT OF REACH OF CHILDREN

Persons mixing and loading this product should wear long pants, a long sleeve shirt, gloves, hat and boots because it may cause irritation of nose, throat, eyes and skin. Do not breathe dust. In case of skin contact wash with water. For eyes, wash with water for 15 minutes and get medical attention.

This product is toxic to fish. Discharge from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Dispose of by perforating and either burning in strict accordance with air pollution control laws or by burying in a safe place in non-crop lands away from water supplies.

NET CONTENTS 50 LBS.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not use this product for any uses other than those specified on this label.

CROP	CROP PEST	DOSAGE RATE/ACRE	LIMITATIONS
Grapes (California Only)	Bunch rot, downy mildew	15 - 40 lbs. Begin application when shoots are 0.5-1.5 inches long. Repeat when shoots are 3-5 inches long, when shoots are 8-10 inches long, and at 10-14 day intervals thereafter until fruit is set.	Do not apply after fruit is set.
Grapes (Areas other than California)	Blackrot, downy mildew	15 - 40 lbs. Begin application when shoots are 0.5-1.5 inches long, repeat when shoots are 3-5 inches long, when shoots are 8-10 inches long and repeat at 10-14 day intervals thereafter.	Do not apply within 66 days before harvest.
Cucumbers, Summer squash, Melons (Cantaloupes, Casababs, Crenshaw, Honeydews, Honey Balls, Muskmelons, Persians, Watermelons)	Alternaria leaf spot, anthracnose, cercospora leaf spot, downy mildew, gummy stem blight, scab	15 - 27 lbs. Begin application at 2 leaf stage and repeat at 5-7 day intervals.	Do not apply within 5 days before harvest.
Potatoes	Early and late blights	10 - 20 lbs. Begin application when plants are 4-6 inches high, and repeat at 7-10 day intervals or more frequently under severe disease conditions.	No time limitations.
Potato Seed pieces		1/2 - 1 lb./100 lbs. of whole or cut seed pieces. Apply for thorough coverage, and plant as soon after treatment as possible.	Do not use treated seed pieces for food or feed.

Fusarium seedpiece decay and seedborne common scab

SEED TREATMENT USES

For control of seed and soil borne fungi causing seed decay, seedling blights, covered smut of barley, rye and wheat, seed borne rust (Puccinia Carthami) of safflower, and fungi on tomato seed.

CAUTION: Do not use treated seed for food, feed or oil purposes.

CROP	SLURRY TREATMENT DOSAGE RATE	DRY TREATMENT DOSAGE RATE
Barley	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Field Corn	15/16 - 1 1/4 lbs./bushel	15/16 - 1 1/4 lbs./bushel
Cotton (acid delinted)	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Cotton (reginned)	3 1/4 lbs./bushel	3 1/4 lbs./bushel
Flax	2 1/2 lbs./bushel	1 1/4 - 2 1/2 lbs./bushel
Oats	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Shelled peanuts	1 1/4 - 2 1/2 lbs./bushel	1 1/4 - 2 1/2 lbs./bushel
Rice	1 1/4 - 2 1/2 lbs./100 lbs. seed	1 1/4 - 2 1/2 lbs./100 lbs. seed
Rye	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Safflower	1 1/4 lbs./bushel	1 1/4 lbs./bushel
Sorghum	15/16 - 1 1/4 lbs./bushel	15/16 - 1 1/4 lbs./bushel
Tomatoes	5 lbs./100 lbs. seed	None
Wheat	1 1/4 lbs./bushel	1 1/4 lbs./bushel

Children.

WARNING

Keep this and all contents of this container out of the reach of children. See back of label for complete directions. Do not use if label is missing or torn. Do not use if contents are empty.