

Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Collards, Cucumbers, Endive, Kale, Lettuce, Lima Beans, Musk Melons, Onions, Parsley, Peas, Peppers, Snap Beans, Spinach, Summer Squash, Sweet Corn, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelon, Winter Squash

Sweet Potatoes

FOR FIELD CROP INSECTS

Peas and Lentils

Field Beans (Broad)

European Corn Borers

Corn Rootworm Larvae and European Corn Borers (North Central US only)

East Armyworms, Southwestern Cornstalk Borers

Cutworms

Lesser Cornstalk Borers

Peanuts

Southern Corn Rootworms

Lesser Cornstalk Borers

Sorghum

Cutworms

Lesser Cornstalk Borers

Soybeans

Cutworms

Lesser Cornstalk Borers

Sugar Beets

Sugar Beet Root Maggots

Wireworms

Sugar cane

Sugar cane Borers

Wireworms

Tobacco

Cutworms

White Grubs

Small Corn Weevils

Wireworms

FOR TURF INSECTS

Centipedes

Grasshoppers

Ground Squirrels, Chipmunks, Skunks, Raccoons, and other mammals

Worms

Top 4-8 inches of soil immediately

7-21 lbs. Apply at planting, working in bands 3-6 inches deep over both sides of rows

2-28 lbs. Apply broadcast before planting and work into top 4-8 inches immediately

7-14 lbs. Apply at planting, working in bands 3-6 inches deep over both sides of rows

2-27 lbs. Apply in 10 inch band centered over row at planting

4 1/2-8 1/2 lbs. 1000 linear feet of row. Apply in a band over one or both sides of row after planting, covering and cover treated band with 2-3 inches of soil

7-14 lbs. Apply so granules fall into whorls when 75% of plants show first generation borer feeding. Apply again 7 days later if needed. For second generation borer control, apply when counts of egg masses are 100 per 100 plants.

7-14 lbs. Apply over plants and cover granules falling to ground with 2-3 inches of soil at peak of first generation corn borer egg laying. Repeat 7 days later if needed. For second generation borers, apply when counts of egg masses are 100 per 100 plants.

7-14 lbs. Apply over corn so granules fall into whorls when armyworms first appear. Repeat if necessary.

14-28 lbs. For surface cutworms, apply just before planting or when larvae first appear, and work into 1-2 inches of soil. For subterranean cutworms, apply just before planting and work into top 3-6 inches of soil immediately

8 1/2-17 or 1000 ft. of row. Apply in a 10 inch band centered over row at planting or when crop starts to emerge. Cover treated band with 1-2 inches of soil

Do not apply within 21 days before harvest. Do not forage within 7 days after application. Do not feed hay within 21 days after application.

Same

Same

Same

Same

Same

Do not apply within 21 days before harvest. Do not forage within 7 days after application. Do not feed hay within 21 days after application.

Same

Do not apply within 7 days before harvest

Same

Do not apply within 7 days before harvest

Same

Do not apply this product in sweet beets with seed, apply wet granules

Same

Do not apply within 28 days before harvest. Do not make more than 10 cuts in the growing season. Do not feed hay and do not use for stock within 21 days of last application

Same

Do not apply within 7 days before harvest

Do not apply within 7 days before harvest

Do not apply within 7 days before harvest

Do not apply within 7 days before harvest