

ACCEPTED
 NOV 13 1974
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER-
 ED UNDER NO. 1528-38

RECEIVED
 23 OCT 1974
 REGISTRATION DIVISION
 PESTICIDES, EPA

TAKE TIME  OBSERVE LABEL DIRECTIONS

**LIVESTOCK REMEDY—
 NOT FOR HUMAN USE
 For External Use Only**

DIRECTIONS FOR USE

For Use on Cattle, Sheep, and Horses

To Operate:

Hold container outlet about 3 inches from wound or infested area. Be sure opening is directed toward area to be treated. Press valve down to release spray. Wet thoroughly by spraying directly into and about 1" around wound.

Screw-worm and Blowfly Infestations:

Use particular care to spray into deep pockets caused by screw-worms. Re-treat at 5 to 7 day intervals as necessary.

To prevent Screw-worm and Blowfly Infestations:

Apply directly to all open or fresh wounds resulting from dehorning, castrations, shear cuts, etc. Also treat navels of newborn animals. Freely bleeding wounds may require several applications. Re-treat at 5 to 7 day intervals as needed.

Fleece Worm Infestations:

Spray infested areas liberally, working toward the center of infestation. Shearing or removing wool from the infested area is not necessary. Re-treat at 5 to 7 day intervals as needed.



**LIVESTOCK
 PRESSURIZED SPRAY
 Contains-KORLAN* Insecticide
 WOUND PROTECTANT—
 Kills screwworms, blowfly maggots
 For livestock of all ages, including newborn animals**

ACTIVE INGREDIENTS:

Ronnel /O, O-dimethyl
 O-(2,4,5-trichlorophenyl)
 Phosphorothioate/ 2.5%
 Xylene 0.5%

INERT INGREDIENTS: 97.0%

EPA Reg. No. 1528-38

EPA Est. 1528-NB-1

WARNING

**KEEP OUT OF REACH OF
 CHILDREN**

Read complete precautions on side panel
 Net Weight 4 oz.

Kills Fast — Saves Time

Worms Leave Wound Quickly

Blue color added for marking treated areas

WARNING

Keep out of reach of children. Injurious to eyes. Do not get in eyes. In case of contact with human eyes, flush eyes immediately with plenty of water and get prompt medical attention. Avoid breathing spray mist. Harmful if swallowed. Avoid skin contact. Wash hands thoroughly after using. Avoid contamination of feed and foodstuffs.

WARNING: Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is an antidote. Consult a veterinarian at the first sign of an adverse reaction.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

*Trademark of The Dow Chemical Company

Manufactured by

SHEPARD LABORATORIES

Division of
NEBRASKA PRODUCING & REFINING
 1521 North 11th Street
 Omaha, Nebraska 68110