

NOTICE OF PESTICIDE: ☒ REGISTRATION
☐ REREISTRATION

*(Under the Federal Insecticide Fungicide
and Rodenticide Act as amended.)*

DATE OF ISSUANCE

DATE OF ISSUE
FEB 15 1985

NAME OF PESTICIDE PRODUCT

BEST AVAILABLE COPY

DAY 4

g. Storage stability--It is not acceptable to reference a storage stability study of a product that is more concentrated than your product. Dilute products tend to be less stable. We can accept the cited study as interim data. However, you must either submit a storage stability study for your product or reference a study of a product that more closely resembles your product.

h. Viscosity--Must be expressed in centistokes or centipoise.

2. Add the phrase "EPA Registration No. 1528-44" to your label before you release the product for shipment.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

RD-37:RKing:kes:2/13/85:Kendrick & Co.-898-1270:Del.2/21/85

BEST AVAILABLE COPY

Ready To Use Pyrethrin Fly Spray
From
PYROCID Intermediate 5192

ACTIVE INGREDIENTS

Pyrethrine	0.50%
*Piperonyl butoxide, technical	1.00%
**N-Octyl bicycloheptene dicarboximide	1.67%
INERT INGREDIENTS	96.83%
	100.00%

*Equivalent to .80% (butylcarbityl) (6-propylpiperonyl)
ether and .20% related compounds.
**HGX 264, Insecticide Synergist.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed. Do not contaminate feed, water or mineral supplements with this spray. Do not contaminate milk or milk handling equipment. Wash udder and teats before milking. Avoid overspraying which will result in wetting the animal's hide.

Environmental Hazards

This product is toxic to fish. Do not contaminate any body of water with this product.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place. Keep container closed.

Disposal: Do not re-use empty container. Wrap container and put in trash collection.

To protect dairy cows and cattle from stable, horn, houseflies, horseflies, mosquitoes and gnats: Spray should be applied daily before the morning or evening milking. Apply as a light mist not to exceed two (2) ounces per animal. Apply from a distance of two feet, permitting the spray particles to collect upon the hair coat. Give special attention to shoulder blades, belly and legs. Wash teats and udder with soap and warm water prior to milking. Do not soak the hide.

FEB. 1995

1528-44

BEST AVAILABLE COPY

For Fogging: Use approximately one (1) ounce of material for each 1,000 cubic feet of space. Keep doors and windows closed for at least ten minutes. Sweep up and destroy all fallen flies. Follow recommendations of equipment manufacturer for proper positioning of the fogging machine.

For Calves: Spray very lightly once or twice daily.

Other Livestock, Horses and Swine: Spray very lightly once or twice daily.

To Protect Cattle and Other Livestock (see above) from Attack by Lice and Ticks: The animals should be sprayed at the rate of two (2) ounces per spraying for adult animals the size of a cow. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks. Ticks must be contacted directly with the spray.

To Protect Cattle and Other Livestock (see above) from Attack by Fleas: Spray bedding and walls of pen, particularly cracks and crevices. The livestock should be sprayed daily until infestation is controlled.

To Protect Cattle and Other Livestock (see above) from Attack by Face Flies: Apply spray to the faces of cattle at the rate of one (1) ounce per animal, paying particular attention to the areas near the eyes and nostrils. Wipe on with a heavy cloth or sponge, or spray on with a sprayer that delivers coarse droplets. Avoid spraying directly into the animal's eyes. Treatment should be made daily while the cattle are stanchioned for the morning milking to give protection from face flies.

Net Contents 1 gal.

EPA Reg. No. 1528-
EPA Est. No. 0677-IA01

Shep's American, Inc.
Denison, IA 51442