



FCX 5% CHLORDANE DUST

INGREDIENTS

ACTIVE INGREDIENT.	
Technical Chlordane *	5 %
INERT INGREDIENTS	95 %
TOTAL	100 %

* Equivalent to 3 percent octachloro 4, 7 methanotetrahydroindane and 2 percent related compounds.

EPA Reg. No. 1598-113 EPA EST 1598-SC-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust or vapor. In case of contact with skin, wash with soap and water. Do not contaminate feed and foodstuffs.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by adequate rainfall or by thorough irrigation, and the grass or soil has completely dried. Do not graze or feed or sell for feeding any forage (e.g. hay from ornamental grass turf) from treated areas to livestock and poultry.


This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees exposed to direct treatment and to residues in or on treated areas. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Bury empty containers in a pit on non-crop land in an isolated area well away from water supplies.

See remainder of Caution statement on back panel.

NET WEIGHT Manufactured By 5 Pounds



FCX 5% CHLORDANE DUST

DIRECTIONS FOR USE

ORNAMENTAL TURF To control ants, chiggers, sod webworm, crickets, cutworms, and earwigs, apply 25 to 30 pounds of dust per acre (6 ounces per 500 square feet of area) uniformly over and into the grass. Apply where the insects are first observed, and repeat as necessary, if the insects are determined to have become reestablished.

For fire ants in ornamental turf and lawns, break open hard surface of mound and pour in 2 to 4 cupfuls of dust per mound; sprinkle dust over areas surrounding mounds as for other ants. If fire ants survive, repeat in 2 weeks.

To control white grubs and Japanese beetle larvae, apply 150 pounds of dust per acre (2 pounds per 500 square feet of area) uniformly over and into the grass. Apply in the fall, preferably, or in early spring. Repeat when the insects are determined to have become reestablished in each particular area; this should not be more frequently than annually.

Rake the turf prior to treatment to aid penetration of the chemical into the soil. Apply when rainfall is not expected within 24 to 48 hours. If adequate rainfall then does not occur, sprinkle the lawn heavily 24 to 48 hours after the application to wash the chemical into the soil. The dust must be washed into the soil, either by adequate rainfall or by sprinkler, 24 to 48 hours after application, (not earlier).

TOMATOES, CABBAGE, PEPPERS (Preplanting Soil Treatment Only) - To control cutworms, apply 25 to 30 pounds of dust per acre (6 ounces per 500 square feet of area) uniformly over the surface of the soil prepared for planting and cultivate into the top 3 to 4 inches of soil prior to setting the transplants of the above named crops. If desired, place 2 to 3 teaspoonfuls of dust in a ring around each transplant, but not in direct contact with the plant. Repeat annually at beginning of planting season.

CABBAGE, BROCCOLI, AND CAULIFLOWER (Pre-planting Soil Treatment Only) - To control cabbage root maggot, apply 75 to 100 pounds of dust per acre (1 to 1 1/2 lb. per 500 sq. ft. of area) uniformly over the surface of the soil prepared for planting and cultivate into the top 3 to 4 inches of soil prior to the setting of transplants of the above named crops. Repeat annually at beginning of planting season.

BEANS, PEAS, LIMA BEANS, AND SWEET CORN (Preplanting Soil Treatment Only) To control seed corn maggot, apply 25 to 30 pounds of dust per acre (6 ounces per 500 sq. ft. of area) uniformly over the surface of the soil prepared for planting and cultivate into the top 3 to 4 inches of soil prior to planting seeds of the above named crops. Repeat annually at beginning of planting season.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label. ODF 374

